

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 October 2001 (11.10.2001)

PCT

(10) International Publication Number
WO 01/74827 A1

(51) International Patent Classification⁷: C07F 9/26,
A61K 31/66, 31/70, A61P 35/00, C07F 9/655, 9/6553,
9/24, C07H 19/10

(21) International Application Number: PCT/US01/10163

(22) International Filing Date: 30 March 2001 (30.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/193,918 31 March 2000 (31.03.2000) US

(71) Applicant: PURDUE RESEARCH FOUNDATION
[US/US]; 1291 Cumberland Avenue, West Lafayette, IN
47906 (US).

(72) Inventors: BORCH, Richard, F.; 2013 Durkees Run
Court, Lafayette, IN 47905 (US). TOBIAS, Sandra, C.;
2408 Kestral Boulevard, Apt. E, West Lafayette, IN 47906
(US). GARRIDO-HERNANDEZ, Hugo; 2408 Kestral
Boulevard, Apt. E, West Lafayette, IN 47906 (US).

(74) Agent: LAMMERT, Steven, R.; Barnes & Thornburg, 11
South Meridian Street, Indianapolis, IN 46204 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHOSPHORAMIDATE PRODRUGS

(57) Abstract: Novel phosphoramidate derivatives of hydroxy functional or amino functional compounds, including amino acids, peptides, peptidomimetics and nucleotide analogs, are described. The compounds enable enhanced intracellular delivery of drugs as their corresponding phosphate esters or amides. Described phosphoramidate compounds exhibit antiproliferative activity. Pharmaceutical formulations are provided for treatment of cancers.

WO 01/74827 A1

PHOSPHORAMIDATE PRODRUGS

Field of the Invention

This invention relates to intracellular delivery of phosphate-substituted therapeutic compounds. More particularly, the invention is directed to certain phosphate precursors and methods and intermediates therefor.

Background and Summary of the Invention

Significant research efforts have been directed towards developing systems for enhancing the intracellular delivery of phosphate-substituted compounds, such as biologically active peptides and nucleotides, inasmuch as the high negative charge of these compounds leads to very low cellular permeability. For example, there is much interest in the intracellular delivery of tyrosine phosphate peptides and peptidomimetics for use in blocking the uncontrolled proliferation of tumor cells. In this regard, protein tyrosine kinases are important regulators of cell cycle progression, and represent attractive targets for the rational design of novel anticancer agents. Protein tyrosine kinases are activated upon phosphorylation of specific tyrosine residues of the kinases and the phosphotyrosine residue along with the three adjacent amino acids (pY-E-E-I) serve as a binding site for Src homology 2 (SH2) domains of intracellular signaling proteins. Interaction between an SH2 domain of an intracellular signaling molecule and an activated protein tyrosine kinase is essential for effective cell cycle progression. Thus, molecules that mimic the phosphotyrosine moiety of protein tyrosine kinases and block the binding of protein tyrosine kinases to SH2 domains represent promising antiproliferative agents for the treatment of cancers.


The phosphate dianion on tyrosine-phosphorylated protein tyrosine kinases is critical to the kinase-SH2 domain interaction. The crystal structures of a number of ligand-SH2 domains have been reported, the interactions have been modeled, and many different inhibitors have been synthesized. In all of these phosphotyrosine peptides and peptidomimetic inhibitors, the phosphotyrosine residue is involved in a complex network of hydrogen bonding and charge-charge interactions that require extensive interaction of the SH2 domain with the phosphate dianion of the tyrosine phosphate moiety. The importance of this dianionic interaction is illustrated

-2-

by the fact that substitution of the phosphate group with a wide variety of other anionic and neutral hydrogen-bonding substituents leads to a significant loss in binding affinity. Thus, the intracellular delivery of tyrosine phosphate peptides and peptidomimetics where the phosphate is in the form of the dianion is critical.

5 Nucleotide analogs represent another class of promising agents for use in treating disease states caused by uncontrolled viral or cellular replication. Like tyrosine phosphate peptidomimetics, the phosphate dianion of nucleotide analogs is critical to the capacity of these compounds to block viral or cellular replication. The most widely used approach for the use of nucleotide analogs as therapeutic agents is
10 to deliver the nucleoside analog as a prodrug across the cell membrane and to rely on the cellular machinery to link a phosphate group to the purine or pyrimidine base to form the corresponding nucleotide analog. However, many nucleosides with therapeutic promise are not converted to the corresponding nucleotide by existing cellular pathways. Furthermore, tumor cells or virus-infected cells may become
15 resistant to treatment by down-regulating the activity of nucleoside kinases, and, for many previously developed nucleotide prodrugs, the intracellular activation process is so inefficient that effective nucleotide concentrations do not accumulate within the cell.

The general strategy for the intracellular delivery of phosphate-
20 containing compounds utilized in the present invention is outlined in Scheme 1:



25

Scheme 1

Briefly, the phosphate-substituted compound to be delivered intracellularly is synthesized (R with attached phosphate in scheme 1 above) and is linked to a delivery
30 group and a masking group. The masking group consists of a haloalkylamine moiety that is hydrolyzed intracellularly to produce an intermediate that is stable as long as the delivery group remains intact, but is released upon intracellular activation of the

-3-

delivery group. The delivery group may be a group, such as a nitrofuryl group or a perhydrooxazine, that is subject to intracellular hydrolysis to provide an intermediate which undergoes cyclization and P-N bond cleavage (i.e., spontaneous hydrolysis) in the intracellular environment releasing the desired phosphate-containing compound.

5 Systems for the intracellular delivery of alkylating agents and nucleotides that employ chemistry related to that described in scheme 1 have been developed previously. However, previous methods resulted in only about 50% intracellular conversion to the desired nucleotide with the other 50% of the prodrug being "lost" as a biologically inactive solvolysis product. The present invention is
10 directed to a newly discovered modification of this chemistry that extends its applicability both to nucleotide analogs and to the phenol phosphate group that is the essential constituent of tyrosine phosphate peptides and peptidomimetics. The invention is based on the discovery that changing the structure of the masking group from a haloethyl to a halobutyl (or halopentyl) group results in the rapid and
15 quantitative intracellular conversion (i.e., about 100% conversion) to the desired nucleotide or phosphate-containing peptide. The new chemistry of the invention is applicable to nucleotide analogs, phosphotyrosine peptides and peptidomimetics, and to a wide variety of other pharmaceutically significant compounds containing phosphate moieties.


20 The development of protected phosphotyrosine precursor phosphoramidates in accordance with this invention enables facile synthesis of phosphotyrosine peptides and peptidomimetics. Thus, the invention is also directed to protected phosphotyrosine precursors and related protected peptides, for example, with N-BOC and fluorenylmethoxycarbonyl groups.

25 In one embodiment of this invention there is provided a phosphoramidate prodrug compound wherein a biologically active compound having hydroxy functionality is covalently linked through that hydroxy functionality to form a biologically labile phosphoramidate which is converted intracellularly to the corresponding drug phosphate. The biologically labile phosphoramidate group
30 enables more efficient intracellular delivery of the drug substance and serves as a means for achieving biologically significant intracellular concentrations of the drug substance in the form of its corresponding phosphate ester. In another embodiment

-4-

the biologically active compound has amino functionality, and it is linked through that amino functionality to form a biologically labile phosphoramidate which is converted intracellularly to the phosphoramidate (OPO₂NH-Drug).

In another embodiment this invention provides an intermediate
5 halophosphoramidite of the formula



wherein n is 4 or 5, R is lower alkyl or -(CH₂)_nX, X is an electrophilic group capable
10 of being nucleophilically displaced from the carbon atom to which it is bound and the group R₁CH₂- is a biologically label ester forming group, more particularly, an ester forming group that is readily hydrolyzed intracellularly. This compound can be used as an intermediate reagent for preparing the above-described phosphoramidate
15 biologically active compound (Drug-OH) is reacted with that halophosphoramidite intermediate to form the corresponding drug reacted intermediate



20 which is then oxidized to form the phosphoramidate prodrug. The method can be used to form phosphoramidite precursors of intracellular drug phosphates starting with hydroxy functional amino acids having carboxyl and amino functional groups protected with standard protecting groups or from biologically active peptides or peptidomimetics or nucleotide analogs. The phosphoramidate prodrugs in accordance
25 with this invention are converted to the corresponding phosphates following hydrolysis of the R₁CH₂- ester forming group. Subsequent hydrolysis of the cyclized 5- or 6-membered zwitterionic intermediate (formed by cyclization of the group -N(CH₂)_nX) provides the corresponding phosphate as the exclusive phosphorous-containing product. Analogous synthetic procedures can be carried out with active
30 compounds having amino functionality represented generally herein as Drug NH₂.

Detailed Description of the Invention

The present invention is directed to novel prodrug chemistry designed to deliver peptide and peptidomimetic phosphates and nucleotide or nucleotide analog phosphates into cells across cell membranes. In one embodiment of the present invention there is provided an intermediate compound of the formula



wherein

R is C₁-C₄ alkyl or -(CH₂)_nX;

n is 4 or 5;

X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;

halo is chloro, bromo or iodo; and

the group R₁CH₂- is a biologically labile ester forming group.

The electrophilic group X is preferably halo, for example, chloro, bromo, or iodo. However, the nature of that group is not critical provided that it can serve as a good leaving group to enable cyclization and concomitant quaternization of the phosphorous-bound nitrogen atom in vivo following administration of the phosphoramidite prodrugs in accordance with this invention. Other electrophilic leaving groups such as acetate, methane sulphonate, trifloromethane sulfonate, haloacetate and the like can be used.


The term "biologically labile ester forming group" as used in defining this invention refers to those ester forming groups derived from alcohols that form ester derivatives that are stable under drug manufacture and storage conditions but are subject to hydrolysis when exposed to biological conditions in vivo. Preferably, the ester forming groups used in accordance with the present invention exhibit minimal susceptibility to hydrolysis in the body fluids in the extracellular space but exhibit susceptibility to hydrolysis in the intracellular space where ester-degrading reductive conditions are prominent. Thus, in one embodiment of the present invention the biologically labile ester forming group on the phosphoramidate prodrugs of this invention are those ester forming groups that are susceptible to hydrolysis under mild

-6-

reductive conditions which include those groups wherein R_1 is nitroaryl including nitrofuryl, nitrothienyl, nitropyrrolyl, nitroimidazolyl, and the like, indanyl, naphthoquinolyl and perhydrooxazine. The nature of the ester forming group R_1CH_2- is not critical provided that the group exhibits susceptibility to hydrolysis under
 5 intracellular conditions, typically biological conditions that exhibit reductive potential.

The therapeutic compounds that can be derivatized in accordance with this invention to form the present phosphoramidate prodrug compounds include amino acids, necessarily in the form of their doubly protected (both carboxyl
 10 protected and amino protected) form, peptides, proteins, peptidomimetics, nucleotide analogs, and other hydroxy functional drug compounds that can exhibit biological activity as their corresponding phosphates. In one preferred embodiment of this invention, the drug is derivatized to the corresponding phosphoramidate in accordance with this invention is a nucleotide analog, such as fluorodeoxyuridine, which exhibits
 15 anti-tumor activity as its corresponding monophosphate derivative. In another embodiment the drug is an amino acid, a peptide or a peptidomimetic.

The phosphoramidate prodrug compounds in accordance with this invention can be prepared from the corresponding hydroxy functional drug substances by any one of several synthetic routes. In one preferred embodiment, the
 20 phosphoramidate prodrug is prepared by a method comprising the steps of reacting the hydroxy functional drug compound with a compound of a formula



under conditions conducive to the formation of an intermediate compound of the formula



and thereafter oxidizing that intermediate to form the present phosphoramidate prodrug of the formula



in which formulas R_1 , R , n , and X are as defined above. The conditions for forming
 30 the intermediate compound of the formula



-7-

can be varied widely, however, it is preferred that the reaction be carried out at low temperature in the presence of an acid scavenger, such as a tertiary amine base. In one embodiment the reaction is carried out at low temperature (about -70°C to about -10°C using dry pyridine as the reaction medium. The oxidation of the intermediate compound of the formula



to the present phosphoramidate prodrug compound of the formula



can be carried out as well under a wide variety of mild oxidizing conditions. The nature of the oxidant is not critical provided that it is capable of affecting the required oxidation of the phosphorous atom to the +5 oxidation state. Examples of oxidizing agents include peracids, peroxides, hydroperoxides and the like. The oxidation is typically carried out at temperatures less than about 0°C, although such is not critical.

The intermediate halo phosphoramidite of the formula



is typically prepared by reacting phosphorous trichloride at low temperature with an alcohol of the formula R_1CH_2OH in an inert organic solvent in the presence of a tertiary amine base, and the resulting product is reacted with an amine of the formula $HNR(CH_2)_nX$. Preferably that intermediate is generated immediately prior to its reaction with the hydroxy functional drug compound, typically in the presence of an acid scavenger, to form the present phosphoramidate prodrug.

In one alternate method of preparing the phosphoramidate prodrugs of this invention the hydroxy functional drug compound is reacted with a chlorophosphoramidate intermediate derived from a reaction sequence starting with phosphorous oxytrichloride (as opposed to phosphorous trichloride) using the same general reactants and conditions described above except that the last oxidation step is eliminated.

Each of the foregoing reactions can be carried out in a similar manner using an amino functional drug (Drug NH_2) to provide the corresponding compounds



In another embodiment of the invention a method is provided of preparing a compound of the formula

-8-



comprising the steps of reacting a compound of the formula



with 1) an alcohol of the formula R_fCH_2OH and 2) an amine of the formula

5 $HNR(CH_2)_nX$, each in the presence of an acid scavenger,

wherein in the above formulas

m is 0 or 1;

R is C_1 - C_4 alkyl or $-(CH_2)_nX$;

n is 4 or 5;

10 X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;

halo is chloro, bromo or iodo; and

the group R_fCH_2- is a biologically labile ester forming group.

15 In yet another embodiment of the invention a method is provided of preparing a phosphoramidate prodrug of the formula



for enhanced intracellular drug delivery of a compound of the general formula $Drug-OPO_3$. The method comprises the steps of reacting a therapeutic compound of the

20 formula $Drug-OH$ with a compound of the formula



under conditions conducive to the formation of the prodrug

wherein in the above formulas

R is C_1 - C_4 alkyl or $-(CH_2)_nX$;

25 n is 4 or 5;

X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;

halo is chloro, bromo or iodo; and


the group R_fCH_2- is a biologically labile ester forming group.

30

As noted above the chemistry described herein can be applied using the same or similar reaction conditions to the preparation of prodrug phosphoramidates of the formula



5 by reaction of the intermediate compound




with a biologically active compound of the general formula Drug-NH₂ wherein R₁, halo, R, n and X are as defined hereinabove. This chemistry provides a means for intracellular delivery of therapeutic compounds of the formula Drug-NHPO₃. Thus,
10 for example, 5'-amino nucleotides or nucleotide analogs can be derivatized as the corresponding prodrug phosphoramidates for efficient intracellular delivery of the corresponding 5'-NHPO₃ derivatives.

The phosphoramidite compounds of the present invention are either themselves prodrugs or, for example, in the instance of tyrosine phosphoramidates,
15 they can be used in the synthesis of peptide or peptidomimetic prodrugs. It will be recognized by those of ordinary skill in the art that the chemistry involved in the synthesis of the phosphoramidites of the present invention is such that the hydroxy, carboxy, thiol, and amine functional groups on the starting drug compound that are not intended to be involved in the chemistry of the phosphoramidate synthesis, should
20 be protected with an art-recognized protecting group. The literature is replete with reference to such protecting groups, their application to protect various functional groups and the conditions for their removal. The use of such protecting groups is well known to those ordinary skill in the art and their use in implementing the synthesis of the present phosphoramidite prodrugs and intermediates therefor form no part of the
25 present invention.

The phosphoramidite prodrugs prepared in accordance with the present invention can be formulated and used in the same or similar dosage forms as the underivatized drug compounds using art-recognized techniques.

In one embodiment of the invention a reaction as outlined in scheme 2
30 is performed:

-10-




Scheme 2

- 5 In this reaction, prodrug 3A was synthesized by sequential addition of nitrofuryl alcohol and p-cresol to phosphoramidic dichloride 1. Prodrug 3 was then activated by reduction with sodium dithionite in ethanol/aqueous buffer, which reduces the nitrofuryl group and leads to the expulsion of phosphoramidate anion 4A. Subsequent reaction of this intermediate was monitored by ³¹P nmr. Intermediate 4A underwent
- 10 smooth cyclization to the 5-membered zwitterionic intermediate, and attack by water gave the cresol phosphate as the exclusive phosphorus-containing product by nmr.

-11-

An alternate embodiment of the invention is illustrated in scheme 3:




Scheme 3

This reaction is directed toward the synthesis of phosphoramidate 11A and involved the preparation of the nitrofuryl phosphoramidate of p-hydroxybenzoic acid 9A and coupling of this product with the benzylamine 10A, prepared from 5-acetylsalicylamide. Briefly, to synthesize the benzoic acid derivative 9A, p-Hydroxybenzoic acid was converted to its allyl ester 6A by reaction of the potassium salt with allylbromide. The allyl ester was reacted with POCl_3 and the intermediate phosphoryl dichloride was treated with methyl chlorobutylamine in the presence of triethylamine to give phosphoramidic chloride 7A. Reaction of 7A with the lithium salt of 5-nitro-2-hydroxymethylfuran afforded the phosphoramidate 8A. Deprotection of the allyl ester was accomplished by reaction with $\text{Pd}(\text{Ph}_3\text{P})_4$ /sodium p-toluenesulfonate to give the benzoic acid phosphoramidate 9A. Reaction of 9A with isobutyl chloroformate in the presence of triethylamine, followed by addition of benzylamine 10A afforded the desired peptidomimetic prodrug 11A. In order to

-12-

confirm that the activation chemistry of 11A to 12A would proceed as expected, 11A was reduced with sodium dithionite in ethanol/aqueous buffer and the reaction monitored by ^{31}P nmr. Conversion of 11 to the phosphoramidate anion was complete within minutes; the phosphoramidate anion was smoothly and exclusively converted to the phosphate 12A with a half-time of 21 minutes at 37°C. The phosphate 12A corresponding to activation of 11A has been synthesized and shown to bind to the pp60^{src} SH2 domain with a binding affinity of 5.6 μM .

In yet another embodiment of the invention, a different intermediate was used for incorporation into the series of reactions depicted in Scheme 3 to form a tyrosine phosphate phosphoramidate. The intermediate 17A is shown below along with the products that result when the reactions depicted in Scheme 3 are performed:




Scheme 3

The intermediate 17A was prepared as described for compound 9A in scheme 3, except that 2-hydroxymethyl-1,4-dimethoxynaphthalene was used instead of 5-nitro-2-hydroxymethylfuran in the third step of the reaction series. Compound 17A was then coupled with amine 10A (Scheme 3) to give the amide 18A using new conditions that generate a high yield of purified product in minutes. Finally, the dimethoxynaphthalene 18A is converted smoothly to the naphthoquinone 19A by oxidation with ceric ammonium nitrate.

-13-

In still another embodiment of the invention, masked phosphotyrosine-containing peptides are synthesized as depicted in scheme 4:



Scheme 4

N-BOC-tyrosine potassium salt was reacted with allyl bromide to give BOC-tyrosine allyl ester 13A. The ester was converted to its cesium phenoxide salt and reacted with N-methyl-N-chlorobutyl phosphoramidic dichloride to give the adduct 14A. Finally, the phosphoramidic chloride was reacted with the lithium salt of 5-nitro-2-hydroxymethylfuran to give the tyrosine phosphoramidate ester 15A. Removal of the allyl protecting group gave the BOC-protected tyrosine phosphoramidate 16A in excellent overall yield. This analog should be suitable for coupling to a variety of amines and will also be valuable as a reagent in automated peptide synthesis.

15

20


The synthesis reaction depicted in scheme 4 has been extended to the development of protected tyrosine phosphoramidates containing the fluorenylmethoxycarbonyl (FMOC) protecting group. FMOC-protected amino acids are widely used in peptide synthesis and offer the advantage that the protecting group can be removed rapidly and cleanly under base-catalyzed conditions. The FMOC analog of tyrosine phosphoramidate 16A, compound 20A above (see Scheme 4) was prepared by a series of reactions similar to those outlined in Scheme 4. The FMOC protecting group is rapidly and cleanly removed by reaction with diazabicycloundecane in dichloromethane for 5 minutes. Finally, coupling reactions of 16A and 20A with methyl glycinate were carried out to demonstrate that these compounds can be used in peptide synthesis reactions.

25

30

-14-

In another embodiment of the present invention, synthesis of the nitrofuryl chlorobutyl nucleoside 3 was carried out in one operation without isolation of intermediates as shown in scheme 5:



Scheme 5

15

Phosphorus trichloride in methylene chloride was reacted with one equivalent of the nitrofuryl alcohol at -78°C and the intermediate dichloride treated with the hydrochloride salt of N-methyl-N-chlorobutylamine to generate the monochloride intermediate A. A solution of 5-fluoro-2'-deoxyriidine in pyridine was added to this intermediate. The resulting phosphoramidate was oxidized with t-butyl hydroperoxide to give the nucleoside phosphoramidate 3. Yields in this synthesis are low and variable (20-60%).

20

In an alternate embodiment of the invention, the analogous thymidine compounds were prepared (i.e., 3 where $\text{F} = \text{CH}_3$). The thymidine phosphoramidate was activated by reduction of the nitrofuryl group with sodium dithionite in aqueous buffer/acetonitrile at 37°C , and the reaction was monitored by ^{31}P nmr. Within ten minutes the resonance for the phosphoramidate anion had disappeared and a new peak corresponding to thymidine-5'-phosphate appeared as the exclusive phosphorus-containing product by ^{31}P nmr. Nucleoside phosphoramidate 3 was then evaluated for its ability to inhibit proliferation of L1210 leukemia cells *in vitro*. Cells were exposed to drug for 2, 8, 24, and 48 hours; the results are summarized below:

25

30

-15-

5


Compound	L1210 IC ₅₀ nM			
	2 hr	8 hr	24 hr	48 hr
3	112	40	6.0	2
5-FU	2742	1390	495	265

The chlorobutyl phosphoramidate **3** is significantly more potent than the analogous bromoethyl compound and is one of the most potent nucleotide prodrugs that we have prepared.

In a related embodiment of the invention, intermediate **4** was incorporated into the above-described reaction scheme and the new drug **5** was produced:

15

20



Scheme 6

The dimethoxynaphthalene analog **4** is synthesized in a manner analogous to that outlined in scheme 5, except that 2-hydroxymethyl-1,4-dimethoxynaphthalene is used instead of 5-nitro-2-hydroxymethylfuran in the first step. The dimethoxynaphthalene **4** is converted smoothly to the naphthoquinone **5** by oxidation with ceric ammonium nitrate. Compound **5** was evaluated an inhibitor of L1210 leukemia cell proliferation in comparison to compound **3** above:

30

-16-

L1210 IC₅₀, nM

	<u>Cpd</u>	<u>2 hr</u>	<u>8 hr</u>	<u>24 hr</u>	<u>48 hr</u>
5	3	116	47	7.2	2.3
	5	44	16	3.1	0.63
	5-FU	2500	1010	360	200

- 10 It is apparent that both compounds are highly potent inhibitors of L1210 cell proliferation, and compound 5 is approximately 3-fold more potent than compound 3 in this assay.

15

20

-17-

In one aspect of the invention the phosphoramidate prodrug is a compound of the formula



formed using a hydroxy or amino functional peptidomimetic (Drug ZH) of the
5 formula:




wherein Z is O or N;

q and k are independently 1 or 0, and

B is H, amino, protected amino, or C₁-C₄ alkanoylamino.


One group of such compounds are benzamido compounds wherein q and k are each 0,
10 for example:




-18-

Another group of such compounds are phenylacetyl compounds wherein q is 1 and k is 0, for example:

5



40




41


10

Still another group of peptidomimetics (Drug ZH) are tyrosyl derivatives wherein q is 1 and k is 1, for example:

15



38



39

Several of such compounds have been found to exhibit significant inhibition of luciferase expression in transfected J77 cells, predictive of antiproliferative activity.

20

Several phosphoramidate compounds prepared in accordance with this invention wherein "Drug ZH" is a nucleotide analog have been found to exhibit significant growth inhibition in L1210 mouse leukemia cells. Such compounds are part of one preferred group of antiproliferative agents of this invention. Numerous nucleotide analogs have been reported in the literature to exhibit antiproliferative activity and any of such compounds, including those that can be covalently bound in


25

-19-


the present phosphoramidates through a hydroxy or an amino functional group can be used in the preparation of the present compounds.

In still another embodiment of the invention the phosphoramidate compounds are derivatives of amino acids and amino acid esters of the formula


5



- 10 wherein Z, q, k and B are as defined above and R is hydrogen or a carboxy protecting group, particularly an ester forming group. Such tyrosine or tyrosine analogs are useful for the preparation of peptides or peptidomimetic compounds capable of facile transmembrane delivery as an intracellular corresponding source of the peptide phosphate or peptidomimetic phosphate. Alternatively such compounds can
- 15 themselves provide an intracellular source of the corresponding amino acid or amino



27



28

acid analog. Examples of such compounds are:

- 20 The present invention further provides pharmaceutical formulations comprising an effective amount of prodrug phosphoramidate compounds for treatment of cancers. As used herein, an effective amount of the prodrug compound is defined as the amount of the compound which, upon administration to a patient, inhibits

-20-

growth/proliferation of tumor cells, kills malignant cells, reduces the volume or size of the tumors or eliminates tumors in the treated patient.

The effective amount to be administered to a patient is typically based on body surface area, patient weight, and patient condition. The interrelationship of dosages for animals and humans (based on milligrams per meter squared of body surface) is described by Freireich, E.J., et al., Cancer Chemother. Rep., 50 (4): 219 (1966). Body surface area may be approximately determined from patient height and weight (see e.g., Scientific Tables, Geigy Pharmaceuticals, Ardley, New York, pages 537-538 (1970)). An effective amount of the phosphoramidate compounds of the present invention, can range from about .05 mg/kg to about 100 mg/kg, about 0.25 mg/kg to about 50 mg/kg, and most typically about 0.1 to about 10 mg/kg per dose. Effective doses will also vary, as recognized by those skilled in the art, dependent on route of administration, excipient usage and the possibility of co-usage with other therapeutic treatments, including other chemotherapeutic agents, and radiation therapy.


The pharmaceutical formulation may be administered via the parenteral route, including subcutaneously, intraperitoneally, intramuscularly and intravenously. Examples of parenteral dosage forms include aqueous solutions or suspensions of the active agent in isotonic saline, 5% glucose or other well-known pharmaceutically acceptable liquid carrier. In one aspect of the present pharmaceutical composition, the phosphoramidate compound is dissolved in a saline solution containing 5% of dimethyl sulfoxide and about 10% Cremphor EL (Sigma Chemical Company). Additional solubilizing agents such as cyclodextrins, which can form more soluble complexes with the present compounds, or other solubilizing agents well-known to those familiar with the art, can be utilized as pharmaceutical excipients for delivery of the present compounds for cancer therapy.

Alternatively, the present compounds can be formulated into dosage forms for other routes of administration utilizing well-known methods. The pharmaceutical compositions can be formulated, for example, in dosage forms for oral administration in a capsule, a gel seal or a tablet. Capsules may comprise any well-known pharmaceutically acceptable material such as gelatin or cellulose derivatives. Tablets may be formulated in accordance with conventional procedure by

-21-

compressing mixtures of the active phosphoramidate and solid carriers and lubricants well-known to those familiar with the art. Examples of solid carriers include starch, sugar, bentonite. The compounds of the present invention can also be administered in a form of a hard shell tablet or capsule containing, for example, lactose or mannitol as a binder and conventional fillers and tableting agents.

Experimental Procedures and Examples



10 **5-Fluoro-2'-deoxyuridyl 5-nitrofurfuryl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate (3):**

Phosphorus trichloride (2M in CH₂Cl₂; 0.81 mL; 1.63 mmol) was cooled to -78°C under argon. 5-Nitrofurfuryl alcohol (233 mg; 1.63 mmol) was dissolved in 12 mL of anhydrous CH₂Cl₂, added to cooled PCl₃ followed by dropwise addition of neat and anhydrous *i*Pr₂NEt (0.57 mL; 3.25 mmol). The reaction was stirred between -78°C and -70°C for 15 min. *N*-Methyl-*N*-(4-chlorobutyl)amine hydrochloride (257 mg; 1.63 mmol) was dissolved in 16 mL of anhydrous CH₂Cl₂ added to the reaction mixture followed by dropwise addition of anhydrous and neat *i*Pr₂NEt (1.13 mL; 6.50 mmol), and the reaction mixture was stirred between -78°C and -70°C for 30 min. The reaction mixture was cannulated to a solution of 5-fluoro-2'-deoxyuridine (200 mg; 0.812 mmol) in 2 mL of anhydrous CH₂Cl₂ and 2 mL of anhydrous pyridine precooled to -65°C. The reaction was stirred between -65°C and -60°C for 1 hour. *t*-Butylhydroperoxide (4.6M in decane; 0.44 mL) was added to the reaction mixture that was stirring at -65°C. The temperature was increased to -40°C and stirred from -40°C to 0°C over 30 min. The reaction mixture was ran through a plug of celite and concentrated under reduced pressure. Column

-22-

chromatography of the crude product (1:1 CH₂Cl₂/acetone) afforded 1 (155 mg; 34%) as a yellow foam.


¹H NMR (CDCl₃): d 9.68 (m, 1H); 7.80 and 7.74 (d, 1H, J = 6.22 Hz and 6.41 Hz); 7.30 (d, 1H, J = 3.48 Hz); 6.71 (d, 1H, J = 3.30); 6.22 (m, 1H); 5.03 (m, 2H); 4.52 (m, 1H); 4.24 (m, 2H); 4.07 (m, 1H); 3.57 (m, 2H); 3.07 (m, 2H); 2.68 (d, 3H, J = 10.25 Hz); 2.49 (m, 1H); 2.18 (m, 1H), 1.74 (m, 4H).

³¹P NMR (CDCl₃, TPPO): d -15.59 and -15.78.

HPLC (35:65 CH₃CN/H₂O [0.1%TFA]): 8.667 min, 92%.

ESI MS (high resolution): Calculated for C₁₉H₃₅ClFN₄O₁₀P: m/z 555.1059 (M+H)⁺;

Found: 555.1059.



5-Fluoro-2'-deoxyuridyl 2-(1,4-dimethoxynaphthyl)methyl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate (4):

Phosphoramidate 4 was prepared from phosphorus trichloride (2M in CH₂Cl₂; 0.41 mL; 0.812 mmol), 1,4-dimethoxy-2-hydroxymethylnaphthalene (177 mg, 0.823 mmol), *N*-methyl-*N*-(4-chlorobutyl)amine hydrochloride (128 mg, 0.812 mmol), 5-fluoro-2'-deoxyuridine (100 mg, 0.406 mmol) and *t*-butylhydroperoxide (4.6M in decane; 0.23 mL) as described above for compound 3. Column chromatography of the crude product (1:1 CH₂Cl₂/acetone) afforded 4 (134 mg; 52%) as a light orange foam.

¹H NMR (CDCl₃): d 8.22 (dd, 1H); 8.04 (dd, 1H); 7.74 and 7.66 (d, 1H, J = 6.23 & 6.39); 7.55 (m, 2H); 6.83 (s, 1H); 6.16 (m, 1H); 5.26 (m, 2H); 4.51 (m, 1H); 4.23 (m, 2H); 4.05 (m, 1H); 3.99 (s, 3H); 3.93 (s, 3H); 3.50 (m, 2H); 3.08 (m, 2H); 2.65 (d, 3H, J = 9.70 Hz); 2.42 (m, 2H); 1.66 (m, 4H).


³¹P NMR (CDCl₃, TPPO): d -12.82 and -12.90

-23-

HPLC (50:50 CH₃CN/H₂O [0.1%TFA]): 7.017 min; 95%.

FAB MS (high resolution): Calculated for C₂₇H₃₄ClFN₃O₄P: m/z 630.1784 (M+H)⁺; Found: 630.1760.

5



10

5-Fluoro-2'-deoxyuridyl 2-(1,4-naphthoquinonyl)methyl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate (5):

15

Ceric ammonium nitrate (225 mg, 0.41 mmol) in water (3 mL) was added dropwise over 15 min to a solution of **4** (100 mg, 0.16 mmol) in CH₃CN (3 mL). The reaction was stirred at room temperature for 1 hour and extracted with CHCl₃ (3x). Combined the organic layers and dried over Na₂SO₄, concentrated under reduced pressure. Purified by silica gel chromatography (5% MeOH:CHCl₃) to give **5** (87.6 mg, 93%) as a yellow foam.

20

¹H NMR (CDCl₃): δ 8.10 (m, 2H); 7.77 (m, 2H); 7.72 (m, 1H); 7.03 (s, 1H); 6.18 (m, 1H); 5.02 (m, 2H); 4.58 (m, 1H); 4.29 (m, 2H); 4.05 (m, 1H); 3.57 (m, 2H); 2.73 (d, 3H, J = 10.07 Hz); 2.47 (m, 1H); 2.31 (m, 1H); 1.75 (m, 4H).

25


³¹P NMR (CDCl₃, TPPO): δ -12.67 and -12.99.

HPLC (50:50 CH₃CN/H₂O [0.1%TFA]): 4.650 min; 100%.

FAB MS (high resolution): Calculated for C₂₅H₂₈ClFN₃O₅P: m/z 600.1314 (M+H)⁺; Found: 600.1327.

30

-24-



**5-Fluoro-2'-deoxyuridyl 2-(5-methoxy-*N*-methylindoloquinone)methyl
N-methyl-*N*-(4-chlorobutyl) phosphoramidate (6):**

5 Phosphorus trichloride (2M in CH_2Cl_2 ; 0.13 mL; 0.258 mmol) was cooled to -60°C under argon. 2-Hydroxymethyl-5-methoxy-*N*-methylindoloquinone (57 mg; .258 mmol) was dissolved in 6 mL of anhydrous CH_2Cl_2 plus 1 mL of anhydrous CH_3CN and added to precooled PCl_3 followed by the dropwise addition of

10 neat i - Pr_2NEt (0.07 mL; 0.386 mmol). The reaction was stirred at -60°C for 25 min. *N*-Methyl-*N*-(4-chlorobutyl)amine hydrochloride (40.7 mg; 0.258 mmol) was dissolved in 1 mL of anhydrous CH_2Cl_2 added to the reaction mixture followed by the dropwise addition of neat i - Pr_2NEt (0.14 mL; 0.773 mmol), and stirred at -60°C for 25 min. 5-Fluoro-2'-deoxyuridine (31.7 mg; 0.129 mmol) was co-evaporated with

15 anhydrous pyridine, dissolved in 2 mL of anhydrous pyridine and cooled to -45°C . This solution of pyridine was titrated with the previously made reaction mixture until FUDR disappeared. The disappearance of FUDR was monitored by TLC using 30% methanol in chloroform. After 1 hour, the reaction was oxidized with t -butylhydroperoxide (4.6 M in decane; 0.06 mL) at -40°C and stirred from -40°C to

20 0°C over 30 min. Saturated NH_4Cl (3 mL) was added, separated the layers and the aqueous layer was extracted with CHCl_3 (5 X 5mL). Combined organic layers and dried over Na_2SO_4 and concentrated under reduced pressure. The crude reaction mixture was passed through a plug of silica gel using 15% methanol in chloroform. Further purification by chromatography on silica gel (1:9 MeOH/ CHCl_3) afforded 6

25 (33 mg; 41%) as a yellow-orange foam.

^1H NMR (CDCl_3): δ 8.99 (br, 1H); 7.62 (m, 1H); 6.62 (d, 1H, $J = 3.76$ Hz); 6.11 (1H, $J = 4.30$); 5.64 (s, 1H); 4.95 (m, 2H); 4.46 (m, 1H); 4.13 (m, 2H); 3.95 (d, 3H, $J = 4.03$

Hz); 3.86 (m, 1H); 3.76 (s, 3H); 3.47 (m, 2H); 3.95 (m, 2H); 2.58 (m, 3H); 2.41 (m, 1H); 2.11 (m, 1H); 1.65 (m, 4H).

³¹P NMR (CDCl₃, TPPO): δ -14.65 and -14.71.


HPLC (Gradient 0% to 70% CH₃CN/H₂O [0.1%TFA] over 30 min): 20.32 min;

5 80%.

FAB MS (high resolution): Calculated for C₂₅H₃₁ClFN₄O₁₀P: m/z 633.1529 (M+H)⁺;

Found: 633.1506.

10



***N*-allyloxycarbonyl Arabinosylcytosine (7):**

Diallyl pyrocarbonate (86.8 mg, 0.452 mmol) was diluted in 1,4-dioxane (5 mL) and added to a solution of arabinosylcytosine (100 mg, 1.61 mmol) dissolved in ddH₂O (1 mL) at room temperature. After the reaction mixture was refluxed for two hours in an oil bath, the solvent was removed under reduced pressure and co-evaporated the white residue with anhydrous pyridine (3 x 5mL). Purified by silica gel chromatography (15% MeOH in CHCl₃) to give 7 (91 mg, 76% based on recovered starting material) as a white foam.


20

25

¹H NMR (DMSO-d₆): δ 10.75 (br, 1H); 8.04 (d, 1H, J = 7.51 Hz); 7.00 (d, 1H, J = 7.51 Hz); 6.04 (d, 1H, J = 3.85 Hz); 5.95 (m, 1H); 5.46 (m, 1H); 5.39 & 5.25 (dd, 1H, J = 1.47 & 35.34 Hz); 5.32 & 5.21 (dd, 1H, J = 1.1 & 23.32 Hz), 5.11 (m, 1H); 4.62 (d, 2H, J = 5.31 Hz); 4.04 (m, 1H); 3.91 (m, 1H); 3.81 (m, 1H); 3.60 (m, 2 H).

30 HPLC (Gradient 0% to 70% CH₃CN/H₂O [0.1%TFA] over 30 min): 12.18 min; 99%. ESI MS (low resolution): (M+H) = 328 m/z; MW = 327 for C₁₃H₁₇N₃O₇.

-26-



5 *N*-allyloxycarbonylarabinosylectosyl 5-nitrofurfuryl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate (**8**):

Phosphoramidate **8** was prepared from phosphorus trichloride (2M in CH₂Cl₂; 1.14 mL; 2.28 mmol), 5-nitrofurfuryl alcohol (326 mg, 2.28 mmol), *N*-methyl-*N*-(4-chlorobutylamine hydrochloride (360 mg, 2.28 mmol), *N*-allyloxycarbonyl arabinosylectosine (240 mg, 0.733 mmol) and *t*-butylhydroperoxide (4.6 M in decane; 0.48 mL) as described above for compound **6**. Column chromatography of the crude product (10% methanol in CHCl₃) afforded **8** (227 mg; 50%) as a light yellow foam.


¹H NMR (CDCl₃): δ 8.23 (d, 1H, J = 7.33 Hz); 7.23 (d, 1H, J = 1.33 Hz); 6.69 (m, 1H); 6.15 (1H); 5.91 (m, 1H); 5.34 (m, 3H); 5.02 (m, 2H); 4.55 (m, 4H); 4.23 (m, 2H); 3.55 (t, 2H, J = 5.68 Hz); 3.03 (m, 2H); 2.67 (d, 3H, J = 10.26 Hz); 1.73 (m, 4H).

³¹P NMR (CDCl₃, TPP): δ -15.54 and -15.76.

HPLC (Gradient 0% to 70% CH₃CN/H₂O [0.1%TFA] over 30 min): 20.01 min; 95%.

ESI MS (high resolution): Calculated for C₂₃H₃₁ClN₅O₁₂P: m/z 636.1474 (M+H)⁺; Found: 636.1474.


-27-



Arabinosylcytosyl 5-nitrofurfuryl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate:

- 5 Tetrakis(triphenyl phosphine) palladium (3.45 mg, 29.9 mmol) was added to a solution of phosphoramidate 8 (38 mg, 59.8 mmol) in THF (400 mL), followed by the addition of *p*-toluenesulfonic acid (11.71 mg, 65.7 mmol) in ddH₂O (240 mL). The reaction mixture was allowed to stir at room temperature for 2.5 hours, after which time it was ran through a plug of silica gel (20% methanol in
- 10 chloroform) to give a light yellow foam (25.1 mg, 76%).
- ¹H NMR (CD₃OD): δ 7.85 (m, 2H); 7.49 (m, 1H); 6.89 (m, 1H); 6.57 (m, 1H); 5.95 (m, 1H); 5.11 (m, 2H); 4.19 (m, 5H); 3.66 (m, 2 H); 3.11 (m, 2H); 2.72 (m, 3H); 1.79 (m, 4H).
- ³¹P NMR (CD₃OD, TPPO): δ -14.18 and -14.37
- 15 **HPLC** (Gradient 0% to 70% CH₃CN/H₂O [0.1%TFA] over 30 min): 21.43 min; 95%.
- ESI MS** (high resolution): Calculated for C₁₉H₂₇ClN₅O₁₀P: *m/z* 552.1262 (M+H)⁺; Found: 552.1251.

-28-



2',3'-Dideoxy-2',3'-didehydrothymidyl 5-nitrofurfuryl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate (10):

5

Phosphoramidate **10** was prepared from phosphorus trichloride (2M in CH₂Cl₂; 0.45 mL; 0.892 mmol); 5-nitrofurfuryl alcohol (128 mg, 0.892 mmol); *N*-methyl-*N*-(4-chlorobutylamine hydrochloride (141 mg, 0.892 mmol); 2',3'-dideoxy-2',3'-didehydrothymidyl (100 mg, 0.446 mmol) and *t*-butylhydroperoxide (4.6M in decane; 0.45 mL) as described above for compound **6**. Column chromatography of the crude product (5% methanol in CHCl₃) afforded **10** (80.5 mg; 48%, based on recovered starting material) as a white foam.

¹H NMR (CDCl₃): δ 8.19 (s, 1H); 7.29 (m, 1H); 7.19 (s, 1H); 6.98 (m, 1H); 6.66 (m, 1H); 6.34 (m, 1H); 5.91 (m, 1H); 4.99 (d, 3H, J = 9.97 Hz); 4.16 (m, 2H); 3.56 (t, 2H, J = 5.59 & 6.13 Hz); 3.06 (m, 2H); 2.63 (m, 3H); 1.87 (2s, 3H); 1.71 (m, 4H).


³¹P NMR (CDCl₃, TPPO): δ -13.58 and -13.99.

HPLC (Gradient 0% to 70% CH₃CN/H₂O [0.1%TFA] over 30 min): 22.75 & 23.18 min; 90%.

ESI MS (high resolution): Calculated for C₂₀H₂₆ClN₄O₉P: m/z 533.1358 (M+H)⁺;

20 Found: 533.1350.


-29-



2',3'-Dideoxy-2',3'-didehydrothymidyl 2-(1,4-dimethoxynaphthyl)methyl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate (11):

- 5 Phosphoramidate 11 was prepared from phosphorus trichloride (2M in CH_2Cl_2 ; 0.45 mL; 0.892 mmol); 1,4-dimethoxy-2-hydroxymethylnaphthalene (195 mg, 0.892 mmol); *N*-methyl-*N*-(4-chlorobutyl)amine hydrochloride (141 mg, 0.892 mmol); 2',3'-dideoxy-2',3'-didehydrothymidyl (100 mg, 0.446 mmol) and *t*-butylhydroperoxide (4.6M in decane; 0.45 mL) as described above for compound 6.
- 10 Column chromatography of the crude product (5% methanol in CHCl_3) afforded 11 (121 mg; 56%, based on recovered starting material) as a white foam.
- ^1H NMR (CDCl_3): δ 8.24 (d, 1H, $J = 8.06$ Hz); 8.05 (m, 2H); 7.54 (m, 2H); 7.19 (s, 1H); 6.98 (s, 1H); 6.84 (s, 1H); 6.31 (m, 1H); 5.58 (m, 1H); 5.24 (d, 2H, $J = 7.69$ Hz); 4.98 (m, 1H); 4.15 (m, 2H); 4.98 (m, 6H); 3.50 (t, 2H, $J = 5.86$ & 6.31 Hz); 3.04, m, 15 2H); 2.61 (m, 3H); 1.86 (s, 3H); 1.71 (m, 4H).
- ^{31}P NMR (CDCl_3 , TPPO): δ -13.44 and -13.88.
- HPLC (Gradient 0% to 70% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ [0.1%TFA] over 30 min): 28.08 & 28.30 min; 87%.
- ESI MS (low resolution): Calculated for $\text{C}_{28}\text{H}_{35}\text{ClN}_3\text{O}_8\text{P}$: 608 ($\text{M}+\text{H}$) $^+$; Found: 630 20 (Na adduct).

-30-



10 **2',3'-Dideoxy-2',3'-didehydrothymidyl 2-(1,4-naphthoquinonyl)methyl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidate (12):**

Ceric ammonium nitrate (225 mg, 0.411 mmol) in water (3 mL) was added dropwise over 15 min to a solution of **11** (100 mg, 0.164 mmol) in CH₃CN (3 mL). The reaction was stirred at room temperature for 1 hour and extracted with CHCl₃ (3x). Combined the organic layers and dried over Na₂SO₄, concentrated under reduced pressure. Purified by silica gel chromatography (5% MeOH:CHCl₃) to give **12** (78 mg, 82%) as a yellow foam.

¹H NMR (CDCl₃): δ 8.09 (m, 2H); 8.01 (s, 1H); 7.79 (m, 2H); 7.17 (s, 1H); 6.99 (m, 2H); 6.35 (m, 1H); 5.93 (m, 1H); 5.00 (m, 3H); 4.24 (m, 2H); 3.56 (t, 2H, J = 5.86 & 6.31 Hz); 3.11 (m, 2H); 2.71 (d, 3H, J = 10.07 Hz); 1.90 (s, 3H); 1.74 (m, 4H).


20 ³¹P NMR (CDCl₃, TPPO): δ -13.49 and -13.84.

HPLC (Gradient 0% to 70% CH₃CN/H₂O [0.1%TFA] over 30 min): 24.82 & 25.02; 95%.

FAB MS (high resolution): Calculated for C₂₅H₂₉ClN₃O₈P: m/z 600.1279 (Na⁺ adduct); Found: 600.1297 (Na⁺ adduct).

25

-31-




5

***N*-methyl-*N*-(4-bromobutyl)amine hydrobromide:**

Hydrobromic acid (20 mL, 0.38 mmol, 48% by wt) was added slowly with stirring to *N*-methyl-*N*-butan-4-ol (4.0 g, 0.39 mmol) at 0°C. The reaction mixture was heated to reflux for 2 hours. A distillation apparatus was attached and 10 mL of distillate was collected, an additional 10 mL (0.18 mmol) of 48% HBr was added. After refluxing for 4 hours, 10 mL were distilled off, an additional 10 mL of 48% HBr (0.18 mmol) were added and the reaction was refluxed overnight. The reaction mixture was then distilled (~20 mL of distillate) and the still pot residue was poured into acetone at -78°C. A white precipitate formed and was collected by filtration (4.24 g, 44%).

¹H NMR: CDCl₃: δ 9.08 (s, 1H); 3.46 (t, 2H, J = 5.95 Hz); 3.04 (m, 2H); 2.72 (t, 3H, J = 5.67 Hz); 2.05 (m, 4H); 1.70 (s, 1H).
D₂O: 2.84 (t, 2H, J = 6.13 Hz); 2.39 (t, 2H, J = 7.32 Hz); 2.04 (s, 3H); 1.23 (m, 4H).

20

***N*-methyl-*N*-(4-chlorobutyl)amine hydrochloride:**

To a solution of *N*-methyl-*N*-butan-4-ol (2.0 g, 19.38 mmol) in 10 mL of CH₂Cl₂ stirring at room temperature, HCl gas was bubbled until it turned litmus paper red (pH=2). The reaction mixture was cooled to 0°C, thionyl chloride (1.41 mL, 19.38 mmol) was added dropwise and the reaction was stirred at room temperature overnight. The solvent was removed under reduced pressure to give a white solid (2.9 g, 95%).


¹H NMR: CDCl₃: δ 8.79 (s, 1H); 3.59 (t, 2H, J = 5.95); 2.97 (m, 2H); 2.70 (s, 3H); 2.06 (m, 2H); 1.95 (m, 2H); 1.63 (s, 1H).

30

-32-

D_2O : d 3.50 (m, 2H); 2.75 (m, 2H); 2.55 (s, 3H); 1.70 (m, 4H).

CI MS (low resolution): $(M+H)^+ = 122$ m/z; MW = 121 for $C_5H_{12}NCl$.



***N*-methyl-*N*-(4-chlorobutyl) phosphoramidic dichloride:**


10 *N*-methyl-*N*-(4-chlorobutyl)amine hydrochloride (2.0 g, 12.73 mmol) was dissolved in 20 mL of anhydrous CH_2Cl_2 and cooled to $-40^\circ C$. Phosphorous oxychloride (1.2 mL, 12.73 mmol) was added neat followed by the drop wise addition of triethyl amine (3.6 mL, 25.46 mmol) in 5 mL of CH_2Cl_2 . Warmed reaction to $0^\circ C$ and slowly to room temperature. Allowed reaction to stir at room temperature for 6

15 hours. The reaction mixture was poured over ice, added saturated ammonium chloride and separated layers. Extracted water layer 3x with CH_2Cl_2 , combined organic layers, dried over anhydrous sodium sulfate and concentrated under reduce pressure. Purified by silica gel flash chromatography (3:1 Hex/ EtOAc) to yield a clear oil (2.57 g, 85%).

20 1H NMR ($CDCl_3$): d 3.58 (t, 2H, $J = 5.95$ Hz); 3.29 (m, 2H); 2.84 (d, 3H, $J = 15.93$ Hz); 1.79 (m, 4H).

^{31}P NMR ($CDCl_3$, TPPO): d -6.6.

25



***N*-methyl-*N*-(4-bromobutyl) phosphoramidic dibromide:**


30 Used same precedence as described above (except that reaction was stirred overnight) with *N*-methyl-*N*-(4-bromobutyl)amine hydrobromide (2.0 g, 8.09

-33-

mmol). Purified using flash silica gel chromatography (3:1 Hex/EtOAc) to yield a clear oil (2.0 g, 92%).

^1H NMR (CDCl_3): d 3.46 (t, 2H, $J = 6.05$ Hz); 3.30 (q, 2H); 2.85 (d, 3H, $J = 16.11$ Hz); 1.88 (m, 4H).

5 ^{31}P NMR (CDCl_3 , TPPO): d -6.75.



5-Nitro-2-furylmethyl *N*-methyl-*N*-(4-chlorobutyl) phosphoramidic chloride:


10 Tetrahydrofurfuryl alcohol (240 mg, 1.68 mmol) was dissolved in 10 mL of anhydrous THF and cooled to -78°C . Lithium hexamethyl disilazane (1.84 mL, 1.84 mmol) was added dropwise and the reaction was stirred for 10 min at -78°C . A solution of phosphoramidic dichloride (400 mg, 1.68 mmol) in 10 mL of anhydrous THF was added dropwise to the reaction mixture. Warmed the reaction slowly from -

15 -78°C to -60°C and allowed to stir at -60°C for 1 hour. Quenched with saturated NH_4Cl , separated layers, extracted water layer 3x with EtOAc. Organic layers were combined, dried over anhydrous sodium sulfate and concentrated under reduced pressure. Purified using flash silica gel chromatography (10:1 CH_2Cl_2 /Acetone) to yield a light brown oil (320 mg, 55%).

20 ^1H NMR (CDCl_3): d 7.29, (d, 1H, $J = 3.66$ Hz); 6.72 (d, 1H, $J = 3.66$ Hz); 5.14 (dd, 2H); 3.56 (t, 2H, $J = 5.86$); 3.20 (m, 2H); 2.71 (d, 3H, $J = 13.58$ Hz), 1.79 (m, 4H).

^{31}P NMR (CDCl_3 , TPPO): d -7.88 .

-34-




5-Nitro-2-furylmethyl *N*-methyl-*N*-(4-bromobutyl) phosphoramidic chloride:

Procedure same as described as above starting with nitrofurfuryl

5 alcohol (245 mg, 1.78 mmol) and phosphoramidic dichloride (500 mg, 1.78 mmol). Purified using flash silica gel chromatography (10:1, CH₂Cl₂/Acetone) to yield a light brown oil (360 mg, 52%).

¹H NMR (CDCl₃): δ 7.29, (d, 1H, J = 3.67 Hz); 6.72 (d, 1H, J = 3.66 Hz); 5.14 (dd, 2H); 3.43 (t, 2H, J = 6.32); 3.19 (m, 2H); 2.72 (d, 3H, J = 13.73 Hz), 1.79 (m, 4H).

10 ³¹P NMR (CDCl₃, TPPO): δ -7.88.



Synthesis of Phosphoramidic dichloride.

Diisopropyl amine (8.82 mL, 50.61 mmol) was diluted with 20 mL of dry
 20 dichloromethane and added slowly to a pre-cooled solution of POCl₃ (2.36 mL, 25.31 mmol) and *N*-methyl-4-chlorobutyl amine hydrochloride (4g, 24.31 mmol) in 40 mL of dry dichloromethane at -20 °C. Reaction mixture was stirred for 3.5 hours letting it warm up gradually to 10 °C. Reaction mixture was quenched with saturated ammonium chloride and extracted with dichloromethane. Organic layers were collected, washed with
 25 brine, dried over MgSO₄ and evaporated to dryness to yield a yellow liquid. Crude product was chromatographed using a mixture of 3:1 Hexanes/EtOAc to yield phosphoramidic dichloride as a clear liquid (5.07g, 84% yield).


-35-

¹H NMR (CDCl₃, TMS) d: 3.59 (t, 2H), 3.29 (m, 2H), 2.85 (d, 3H, *j* = 16.11 Hz), 1.80 (m, 4H) ppm.

³¹P NMR (CDCl₃, TPPO) d: -6.63 ppm.

5

10



Synthesis of 1-propenyl benzoate 12a.

Diisopropyl amine (6.31 mL, 36.20 mmol) was added neat to a suspension of 4-hydroxybenzoic acid (5g, 36.20 mmol) in 86 mL of allyl bromide. Reaction mixture was stirred at reflux for 2 hours and allowed to cool to room temperature. Excess allyl bromide was distilled off under reduced pressure and the remaining oil was diluted in EtOAc and washed with water, brine, dried over MgSO₄ and evaporated to dryness to yield a yellow oil. Crude product was chromatographed using a mixture of 10:1 CHCl₃/EtOAc to yield the benzoate ester (5.03g, 78 % yield) as a white solid.

¹H NMR (CDCl₃, TMS) d: 7.99 (d, 2H, *j* = 8.79 Hz), 6.86 (d, 2H, *j* = 8.79 Hz), 6.0 (m, 1H), 5.30 (m, 2H), 4.80 (d, 2H, *j* = 5.68 Hz).


HPLC (50:50 CH₃CN/H₂O-0.1% TFA): 6.17 min (97.2%).

MS (ESI) *m/z* :179 (M+H)

GCMS *m/z*: 179 (M+H).

20

-36-


**Synthesis of 1-propenyl ester 12b.**

Ester **12b** was synthesized as described for **12a** and was obtained as a white solid (78% yield).

5 **¹H NMR (CDCl₃, TMS) d:** 7.77 (d, 2H, *j* = 7.41 Hz), 7.56 (d, 2H, *j* = 7.41 Hz), 7.43-7.28 (m, 4H), 6.96 (d, 2H, *j* = 8.28 Hz), 6.73 (d, 2H, *j* = 8.28 Hz), 5.94-5.80 (m, 1H), 5.35-5.24 (m, 3H), 4.91 (s, 1H), 4.62 (m, 3H), 4.48-4.31 (m, 2H), 4.20 (t, 1H), 3.06 (t, 2H).

HPLC (50:50 CH₃CN/H₂O-0.1% TFA): 11.75 min (99.99%)

10

**Synthesis of 1-propenyl ester 12c.**


15 Ester **12c** was synthesized as described for **12a** and obtained as white solid (80 %yield).

¹H NMR (CDCl₃, TMS) d: 6.99 (d, 2H, *j* = 8.42 Hz), 6.73 (d, 2H, *j* = 8.42 Hz), 5.84 (m, 1H), 5.27 (m, 2H), 4.60 (d, 2H, *j* = 5.86 Hz), 3.02 (m, 1H), 1.42 (s, 9H).

HPLC (30:70 CH₃CN/H₂O-0.1% TFA): 8.95 min (97%).

20

-37-

**Synthesis of 1-propenyl ester 12d.**


Ester **12d** was synthesized as described for **12a** and obtained as a white solid (80% yield)

5 **¹H NMR (CDCl₃, TMS) d:** 7.20-7.09 (m, 4H), 5.99-5.78 (m, 2H), 5.35-5.25 (m, 2H), 4.93-4.85 (m, 1H), 4.61 (d, 2H, *j* = 5.86 Hz), 3.26-3.13 (m, 2H), 2.00 (s, 3H) ppm.

HPLC (30:70, CH₃CN/H₂O-0.1% TFA): 5.63, 5.8 min.

MS (ESI) *m/z*: 286 (M⁺Na).

10



15


Synthesis of 1-propenyl ester 12e.

Ester **12e** was synthesized as described for **12a** and obtained as green oil

¹H NMR (CDCl₃, TMS) d: 7.32-7.19 (m, 4H), 5.98-5.82 (m, 1H), 5.31-5.20 (m, 2H), 4.60 (d, 2H, *j* = 5.68 Hz), 3.64 (s, 2H) ppm.

20 **HPLC (30:70, CH₃CN/H₂O-0.1% TFA):** 4.33 min.

-38-



5


Synthesis of 1-propenyl ester 13.

Allyl bromide (0.99 mL, 11.41 mmol) was added to a suspension of potassium p-aminobenzoate (2g, 11.41 mmol) in 40 mL of DMF. Reaction was stirred overnight and then quenched with water and extracted several times with EtOAc. Organic
 10 extracts were combined, washed with brine, dried over MgSO_4 and evaporated to dryness to yield yellow liquid. Crude product was chromatographed using a mixture of 10:1 $\text{CHCl}_3/\text{EtOAc}$ to yield 13 (2g, 70% yield) as a light yellow solid.

$^1\text{H NMR}$ (CDCl_3 , TMS) δ : 7.99 (d, 2H, $j = 8.7$ Hz), 7.08 (d, 2H, $j = 8.7$ Hz), 6.05-5.97 (m, 1H), 5.51-5.30 (m, 2H), 4.82-4.78 (m, 2H) ppm.

15 **HPLC (gradient 30-100% $\text{CH}_3\text{CN}:\text{H}_2\text{O}$ -0.1% TFA over 35 min): 7.68 min**

MS (ESI) m/z : 178 (M+H), 200 (M+Na).



20

Synthesis of Phosphoramidic monochloride 14a

Diisopropylethyl amine (0.3 mL, 1.68 mmol) was added neat to a pre-cooled solution of phosphoramidic dichloride (0.2 g, 0.84 mmol) and allyl ester 12a (0.15g, 0.84 mmol) in 3 mL of dry DCM at -15°C . Reaction was stirred for 1 hr at -15°C
 25 $^\circ\text{C}$ and 3 hours at -5°C and quenched with saturated ammonium chloride. Layers were separated. Organic layer was then washed with brine, dried over MgSO_4 and evaporated to dryness to yield a yellow liquid. Crude product was chromatographed using a 10:1


-39-

mixture of $\text{CHCl}_3/\text{EtOAc}$ to yield **14a** as a clear liquid.

^1H NMR (CDCl_3 , TMS) δ : 8.10 (d, 2H, $j = 8.33$ Hz), 7.34 (d, 2H, $j = 8.33$ Hz), 6.09-5.98 (m, 1H), 5.45-5.28 (m, 2H), 4.84-4.81 (m, 2H), 3.57 (t, 2H), 3.35-3.17 (m, 2H), 2.84 (d, 3H, $j = 13.55$), 1.82-1.79 (m, 4H) ppm.

5 ^{31}P NMR (TPPO) δ : -13.07 ppm.

HPLC (gradient 30-100%, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA, over 35 min): 16.05 min.



10 Synthesis of phosphoramidic monochloride **14b**.

Phosphoramidic dichloride **11** (0.45g, 1.89 mmol) was dissolved in 2 mL of dry THF and cannulated to a pre-cooled solution of **12b** (0.84g, 1.89 mmol) and LiHMDS (2.08 mmol) in 2 mL of dry THF at -20°C . Reaction was warmed up to 0°C and stirred for 5 hours. Reaction was quenched with saturated ammonium chloride and extracted with ethyl acetate. Organic extracts were combined, washed with brine, dried over MgSO_4 and evaporated to dryness to yield yellow oil. Crude product was chromatographed using a 10:1 mixture of $\text{CHCl}_3/\text{EtOAc}$ to yield **14b** (0.85g, 70% yield) as yellow oil.


^1H NMR (CDCl_3 , TMS) δ : 7.77 (d, 2H, $j = 7.42$ Hz), 7.57 (d, 2H, $j = 7.42$ Hz), 7.44-7.29 (m, 4H), 7.19-7.08 (m, 4H), 5.91-5.81 (m, 1H), 5.34-5.25 (m, 3H), 4.66-4.60 (m, 3H), 4.49-4.33 (m, 2H), 4.21 (t, 1H), 3.58 (t, 2H), 3.27-3.20 (m, 2H), 3.17-3.10 (m, 2H), 2.82 (d, 3H, $j = 13.54$ Hz), 1.81-1.78 (m, 4H) ppm.

^{31}P NMR (CDCl_3 , TPPO) δ : -12.64 ppm.

HPLC (70:30 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% solution of TFA): 6.833 min.

25 MS (ESI) m/z : 645/647 ($\text{M}+\text{H}$), 667/669 ($\text{M}+\text{Na}$).

-40-



Synthesis of phosphoramidic monochloride 14c.


14c was synthesized as described for 14b and obtained as light yellow oil.

- 5 ¹H NMR (CDCl₃, TMS) d: 7.16 (m, 4H), 5.8 (m, 1H), 5.3 (m, 2H), 4.6 (d, 2H, *j* = 5.68 Hz), 3.58 (t, 2H), 3.2 (m, 2H), 3.05 (m, 1H), 2.82 (d, 3H), 1.79 (m, 4H), 1.42 (s, 9H).

³¹P NMR (CDCl₃, TPO) d: -12.5 ppm.

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 9.83 min.

10



Synthesis of phosphoramidic monochloride 14d.


- 15 14d was synthesized in a similar manner as 14a. The reaction mixture was stirred overnight to yield light yellow oil.

¹H NMR (CDCl₃, TMS) d: 7.20-7.09 (m, 4H), 5.99-5.78 (m, 2H), 5.35-5.25 (m, 2H), 4.93-4.85 (m, 1H), 4.61 (d, 2H, *j* = 5.86 Hz), 3.58 (t, 2H), 3.26-3.13 (m, 4H), 2.83 (d, 3H, *j* = 13.55 Hz), 2.00 (s, 3H), 1.96-1.75 (m, 4H) ppm.

³¹P NMR (TPPO) d: 12.71

- 20 MS (ESI) *m/z*: 465/467 (M+H).

-41-



Synthesis of phosphoramidic monochloride 14e.


14e was synthesized in a similar manner as 13a.

¹H NMR (CDCl₃, TMS) d: 7.32-7.19 (m, 4H), 5.98-5.82 (m, 1H), 5.31-5.20 (m, 2H),
 4.60 (d, 2H, *j* = 5.68 Hz), 3.64 (s, 2H), 3.58 (t, 2H), 3.25-3.18 (m, 2H), 2.83 (d, 3H, *j* =
 13.55 Hz), 1.80-1.78 (m, 4H) ppm.

³¹P NMR (TPPO) d: -12.66 ppm.

MS (ESI) *m/z*: 394/396 (M+H).

10



Synthesis of phosphodiamidic monochloride 15.


Diisopropylethyl amine (0.51 mL, 2.94 mmol) was added neat to a mixture of phosphoramidic dichloride 11 (0.7 g, 2.94 mmol) and allyl ester 13 (0.52 g, 2.94 mmol) in 10 mL of dry dichloromethane. Reaction was stirred at reflux for 10 days and allowed to cool down to room temperature, diluted with dichloromethane and quenched with saturated ammonium chloride. Layers were separated; organic layer was washed with brine, dried over MgSO₄ and evaporated to dryness to yield yellow oil. Crude product was chromatographed using a mixture of 10:1 CHCl₃/EtOAc to yield light green oil (0.22 g, 20% yield).

¹H NMR (CDCl₃, TMS) d: 7.99 (d, 2H, *j* = 8.7 Hz), 7.08 (d, 2H, *j* = 8.7 Hz), 6.05-5.97 (m, 1H), 5.57 (d, 1H, *j* = 2.29 Hz), 5.51-5.30 (m, 2H), 4.82-4.78 (m, 2H), 3.54 (t, 2H), 3.36-3.13 (m, 2H), 2.74 (d, 3H, *j* = 14.37 Hz), 1.87-1.64 (m, 4H) ppm.

-42-

³¹P NMR (TPPO) d: -11.28 ppm.HPLC (30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 12.60 min.MS (ESI) *m/z* : 379/381 (M+H), 401/403 (M+Na).

5

**Synthesis of phosphoramidate 16.**


Phosphoramidic monochloride **14a** (0.63g, 1.21 mmol) was dissolved in 2 mL of dry THF and cannulated to a precooled solution of nitrofuryl alcohol (0.19g, 1.33 mmol) and LiHMDS (1.46 mmol) in 2 mL of dry THF at -78° C. Reaction was brought to -40° C and stirred for 5.5 hours. Saturated ammonium chloride was added to the reaction mixture and extracted with ethyl acetate. Organic extracts were combined, washed with brine, dried over MgSO₄ and evaporated to dryness to yield dark oil. Crude product was chromatographed using a mixture of 10:1 CHCl₃/EtOAc to yield phosphoramidate **16** (0.30g, 52% yield) as a dark orange oil.

¹H NMR (CDCl₃, TMS) d: 8.05 (d, 2H, *j* = 8.6 Hz), 7.26 (m, 3H), 6.64 (d, 1H, *j* = 3.48 Hz), 6.05 (m, 1H), 5.31 (m, 2H), 5.09 (d, 2H, *j* = 9.15 Hz), 4.82 (d, 2H, *j* = 5.49 Hz), 3.53 (t, 2H), 3.1 (m, 2H), 2.72 (d, 3H, *j* = 10.25 Hz), 1.67 (m, 4H).

³¹P NMR (TPPO) d: -21.7HPLC (60:40 CH₃CN/H₂O-0.1% solution of TFA): 7.117 min.

20

-43-



5 Synthesis of phosphoramidate 17.

Phosphoramidate 17 was synthesized in a similar manner as described for 16.


¹H NMR (CDCl₃, TMS) d: 8.23 (d, 1H, *j* = 7.14 Hz), 8.07-7.91 (m, 3H), 7.66-7.50 (m, 2H), 7.30 (d, 2H, *j* = 8.7 Hz), 6.80 (s, 1H), 6.19-5.88 (m, 1H), 5.44-5.20 (m, 4H), 4.92-4.79 (m, 2 H), 3.94 (s, 3H), 3.92 (s, 3H), 3.48 (t, 2H), 3.24-2.97 (m, 2H), 2.71 (d, 3H, *j* = 10.25 Hz), 2.84-1.83 (m, 4H) ppm.

³¹P NMR (TPPO) d: -18.94 ppm.

HPLC (70:30 CH₃CN/H₂O-0.01% TFA solution): 7.967 min

MS (ESI) *m/z* : 584/586 (M+Na).

15



Synthesis of phosphoramidate 18.

Phosphoramidate 18 was synthesized in a similar manner as described for

20 16.

¹H NMR (CDCl₃, TMS) d: 7.77 (d, 2H, *j* = 7.42 Hz), 7.57 (d, 2H, *j* = 7.42 Hz), 7.44-7.24 (m, 5H), 7.13-6.93 (m, 4H), 6.59 (d, 1H, *j* = 3.67 Hz), 6.07-5.70 (m, 1H), 5.47-5.23

-44-


(m, 4H), 5.05 (d, 2H, $j = 9.16$ Hz), 4.74-4.61 (m, 3H), 4.56-4.28 (m, 2H), 4.20 (t, 1H), 3.51 (t, 2H), 3.20-2.99 (m, 4H), 2.70 (d, 3H, $j = 10.34$ Hz), 1.82-1.53 (m, 4H) ppm.

^{31}P NMR (TPPO) d: -18.92

HPLC (70:30 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% solution of TFA): 6.12 min.

5

Synthesis of phosphoramidate 19.



10


Phosphoramidate 19 was synthesized in a similar manner as 16.

^1H NMR (CDCl_3 , TMS) d: 7.28 (d, 1H, $j = 3.66$ Hz), 7.16 (m, 4H), 6.63 (d, 1H, $j = 3.66$ Hz), 5.8 (m, 1H), 5.3 (m, 2H), 5.05 (d, 2H, $j = 8.98$ Hz), 4.6 (d, 2H), 3.58 (t, 2H), 3.2 (m, 2H), 3.05 (m, 1H), 2.82 (d, 3H), 1.79 (m, 4H), 1.42 (s, 9H).

^{31}P NMR (TPPO) d: -19.2

15

HPLC (60:40 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA): 8.2 min.



Synthesis of phosphoramidate 20.

Phosphoramidate 20 was synthesized in a similar manner as described for


5 16.

¹H NMR (CDCl₃, TMS) d: 7.28 (d, 1H, *j* = 3.75 Hz), 7.21-6.98 (m, 4H), 6.63 (d, 1H, *j* = 3.75 Hz), 6.05-5.77 (m, 2H), 5.44-5.19 (m, 2H), 5.06 (d, 2H, *j* = 9.15 Hz), 5.01-4.78 (m, 1H), 4.72-4.52 (m, 2H), 3.66-3.42 (m, 2H), 3.23-2.99 (m, 4H), 2.71 (d, 3H, *j* = 10.34 Hz), 2.00 (s, 3H), 1.83-1.60 (m, 4H) ppm.

10 ^{31}P NMR (TPPO) δ : -19.03 ppm.

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 15.70 min.

MS (ESI) m/z : 572/574 (M+H).



15 Synthesis of phosphoramidate 21.

Phosphoramidate **21** was synthesized in a similar manner as described for

16.

¹H NMR (CDCl₃, TMS) d: 7.81 (d, 1H, *j* = 4.12 Hz), 7.22-7.05 (m, 4H), 6.99 (d, 1H, *j* = 4.12 Hz), 6.02-5.72 (m, 2H), 5.45-5.10 (m, 4H), 5.00-4.79 (m, 1H), 4.63-4.49 (m, 2H), 3.53 (t, 2H), 3.23-2.95 (m, 4H), 2.72 (d, 3H, *j* = 10.35 Hz), 2.00 (s, 3H), 1.85-1.58 (m, 4H) ppm.


 ^{31}P NMR (TPPO) d: -19.21 ppm.

20

-46-

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 17.20 min

MS (ESI) *m/z*: 588/590 (M+H), 610/612 (M+Na).



5 Synthesis of phosphoramidate 22.

Phosphoramidate 22 was synthesized in a similar manner as described for 16.


¹H NMR (CDCl₃, TMS) δ: 7.26-7.04 (m, 5H), 6.60 (d, 1H, *j* = 3.58 Hz), 5.99-5.75 (m, 1H), 5.43-5.19 (m, 2H), 5.06 (d, 2H, *j* = 9.16 Hz), 4.72-4.48 (m, 2H), 3.62 (s, 2H), 3.51 (t, 2H), 3.18-2.96 (m, 2H), 2.71 (d, 3H, *j* = 10.26 Hz), 1.77-1.55 (m, 4H) ppm.

³¹P NMR (TPPO) δ: -19.01 ppm.

HPLC (30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 15.50 min.

MS (ESI) *m/z*: 501/503 (M+H), 523/525 (M+Na).

15



Synthesis of phosphoramidate 23.


Phosphoramidate 23 was synthesized in a similar manner as described for 16.

¹H NMR (CDCl₃, TMS) δ: 7.80 (d, 1H, *j* = 4.12 Hz), 7.33-7.09 (m, 4H), 6.99 (d, 1H, *j* = 4.12 Hz), 6.08-5.68 (m, 1H), 5.46-5.07 (m, 4H), 4.78-4.39 (m, 2H), 3.63 (s, 2H), 3.51 (t, 2H), 3.30-2.96 (m, 2H), 2.72 (d, 3H, *j* = 10.34 Hz), 1.87-1.48 (m, 4H) ppm.

³¹P NMR (TPPO) δ: -19.21 ppm.

-47-

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 20.17 min.



Synthesis of phosphoramidate 24.

5 Phosphoramidate **24** was synthesized in a similar manner as described for **16**.


¹H NMR (CDCl₃, TMS) δ: 7.96 (d, 2H, *j* = 8.6 Hz), 7.26 (d, 1H, *j* = 3.67 Hz), 6.99 (d, 2H, *j* = 8.6 Hz), 6.65 (d, 1H, *j* = 3.67 Hz), 6.16-5.84 (m, 1H), 5.51-5.19 (m, 3H), 5.09 (d, 2H, *j* = 9.43 Hz), 4.92-4.73 (m, 2H), 3.52 (t, 2H), 3.22-2.99 (m, 2H), 2.69 (d, 3H, *j* = 10.53 Hz), 1.85-1.61 (m, 4H) ppm.

³¹P NMR (TPPO) δ: -15.60 ppm

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 12.60 min.

MS (ESI) *m/z* : 486/488 (M+H), 508/510 (M+Na).

15



Synthesis of phosphoramidate 25.

TolSO₂Na (0.29g, 1.61 mmol) was dissolved in 2.5 mL of water and added to a solution of phosphoramidate **16** (0.71g, 1.47 mmol) and Pd(PPh₃)₄ (0.08g, 73.41mmol) in 6 mL of THF. Reaction was stirred for 30 minutes at room temperature. Diethyl ether was added to the reaction mixture and washed several times with water. Aqueous extracts were combined, washed with ether, acidified to pH 3 with 2% HCl and extracted with EtOAc. Organic extracts were combined,


-48-

washed with brine, dried over MgSO_4 and evaporated to dryness to yield phosphoramidate **25** (0.51 g, 78% yield) as orange foam.

^1H NMR (CDCl_3 , TMS) δ : 8.08 (d, 2H, $J = 8.6\text{ Hz}$), 7.31 (m, 3H), 6.66 (d, 1H, $J = 3.67\text{ Hz}$), 5.11 (d, 2H, $J = 9.16\text{ Hz}$), 3.53 (t, 2H), 3.13 (m, 2H), 2.74 (d, 3H, $J = 10.44\text{ Hz}$), 1.7 (m, 4H).

^{31}P NMR (TPPO) δ : -20.4

HPLC (60:40 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% solution of TFA): 4.017 min.



Synthesis of phosphoramidate **26**.

Phosphoramidate **26** was synthesized in a similar manner as described for **25**.


^1H NMR (CDCl_3 , TMS) δ : 8.32-8.22 (m, 1H), 8.13-7.98 (m, 3H), 7.64-7.45 (m, 2H), 7.33 (d, 2H, $J = 8.7\text{ Hz}$), 6.81 (s, 1H), 5.35 (d, 2H, $J = 8.05\text{ Hz}$), 3.95 (s, 3H), 3.93 (s, 3H), 3.49 (t, 2H), 3.25-2.97 (m, 2H), 2.73 (d, 3H, $J = 10.25\text{ Hz}$), 1.80-1.52 (m, 4H) ppm.

^{31}P NMR (TPPO) δ : -18.99 ppm

HPLC (gradient 30-100%, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA, over 35 min): 18.60 min.

MS (ESI) m/z : 544/546 ($\text{M}+\text{Na}$).

-49-



Synthesis of Phosphoramidate 27.


Phosphoramidate 27 was synthesized in a similar manner as described

5 for 25.

$^1\text{H NMR (CDCl}_3, \text{TMS) d: 7.77 (d, 2H, } j = 7.23 \text{ Hz), 7.58 (d, 2H, } j = 7.23 \text{ Hz), 7.43-7.23 (m, 5H), 7.18-6.89 (m, 4H), 6.67-6.47 (m, 1H), 5.48-5.30 (m, 1H), 5.18-4.97 (m, 2H), 4.79-4.07 (m, 4H), 3.60-3.37 (m, 2H), 3.27-2.98 (m, 4H), 2.82-2.59 (m, 3H), 1.79-1.56 (m, 4H) \text{ ppm.}$

10 $^{31}\text{P NMR (TPPO) d: -19.44, -19.63 ppm.}$

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 17.93 min.




15 Synthesis of Phosphoramidate 28.

Phosphoramidate 28 was synthesized in a similar manner as described for 25.

$^1\text{H NMR (CDCl}_3, \text{TMS) d: 7.28 (d, 1H, } j = 3.66 \text{ Hz), 7.16 (m, 4H), 6.63 (d, 1H, } j = 3.66 \text{ Hz), 5.05 (d, 2H, } j = 8.98 \text{ Hz), 3.58 (t, 2H), 3.2 (m, 2H), 3.05 (m, 1H), 2.82 (d, 3H), 1.79 (m, 4H), 1.42 (s, 9H).}$

20

-50-

³¹P NMR (TPPO) δ: -19.2 ppm.HPLC (60:40 CH₃CN/H₂O-0.1% TFA): 4.65 min.**5 Synthesis of Phosphoramidate 29.**


Phosphoramidate 29 was synthesized in a similar manner as described for 25.

¹H NMR (CDCl₃, TMS) δ: 7.30-7.25 (m, 1H), 7.23-6.83 (m, 4H), 6.80-6.47 (m, 1H), 5.09 (d, 2H, *j* = 9.25 Hz), 5.01-4.59 (m, 2H), 3.76-3.33 (m, 2H), 3.36-2.98 (m, 4H),

10 2.98-2.55 (m, 3H), 2.14-1.77 (d, 3H), 1.89-1.55 (m, 4H) ppm.

³¹P NMR (TPPO) δ: -19.25, -19.43 ppm.HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 11.68 min.MS (ESI) *m/z*: 554/556 (M+Na), 532/534 (M+H).

15

**Synthesis of Phosphoramidate 30.**

Phosphoramidate 30 was synthesized in a similar manner as described for 25.

¹H NMR (CDCl₃, TMS) δ: 7.87-7.73 (m, 1H), 7.14-7.04 (m, 4H), 7.03-6.95 (m,


-51-

^1H NMR (CDCl₃) δ : 5.34-5.18 (m, 2H), 5.11-4.96 (m, 1H), 4.95-4.81 (m, 1H), 3.60-3.46 (m, 2H), 3.26-3.03 (m, 4H), 2.85-2.67 (m, 3H), 2.05-1.92 (m, 3H), 1.84-1.62 (m, 4H) ppm.

^{31}P NMR (TPPO) δ : -19.41, -19.61 ppm.

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 13.10 min.

5 MS (ESI) m/z : 548/550 (M+H).



Synthesis of Phosphoramidate 31.


Phosphoramidate **31** was synthesized in a similar manner as described

10 for **25**.

^1H NMR (CDCl₃, TMS) δ : 7.26-7.04 (m, 5H), 6.60 (d, 1H, $j = 3.58$ Hz), 5.43-5.19 (m, 2H), 5.06 (d, 2H, $j = 9.16$ Hz), 3.62 (s, 2H), 3.51 (t, 2H), 3.18-2.96 (m, 2H), 2.71 (d, 3H, $j = 10.26$ Hz), 1.77-1.55 (m, 4H) ppm.

^{31}P NMR (TPPO) δ : -19.01 ppm.

15



Synthesis of Phosphoramidate 32.

Phosphoramidate **32** was synthesized in a similar manner as described for

20 **25**.

^1H NMR (CDCl₃, TMS) δ : 7.80 (d, 1H, $j = 4.12$ Hz), 7.33-7.09 (m, 4H), 6.99 (d, 1H,


-52-

$j = 4.12$ Hz), 5.22 (d, 2H, $j = 8.52$ Hz), 3.63 (s, 2H), 3.48 (t, 2H), 3.30-2.96 (m, 2H), 2.73 (d, 3H, $j = 10.25$ Hz), 1.92-1.40 (m, 4H) ppm.

^{31}P NMR (TPPO) d: -19.29 ppm.

HPLC (gradient 30-100%, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA, over 35 min): 15.13 min.

5



Synthesis of Phosphoramidate 33.

10 Phosphoramidate 33 was synthesized in a similar manner as described for 25.


^1H NMR (CDCl_3 , TMS) d: 7.96 (d, 2H, $j = 8.6$ Hz), 7.26 (d, 1H, $j = 3.67$ Hz), 6.99 (d, 2H, $j = 8.6$ Hz), 6.65 (d, 1H, $j = 3.67$ Hz), 6.45-6.25 (m, 1H), 5.09 (d, 2H, $j = 9.43$ Hz), 3.52 (t, 2H), 3.22-2.99 (m, 2H), 2.69 (d, 3H, $j = 10.53$ Hz), 1.85-1.61 (m, 4H) ppm.

15 ^{31}P NMR (TPPO) d: -15.40 ppm

HPLC (gradient 30-100% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA, over 35 min): 9.13, 9.37 min.

MS (ESI) m/z : 444/446 (M-H).

-53-



Synthesis of prodrug 35.

Phosphoramidate **25** (0.105g, 0.23 mmol) was dissolved in 2 mL of dry DMF and cannulated to a mixture of benzylamine **34** (0.0735g, 0.23 mmol), 4-methyl morpholine (0.06 mL, 0.588 mmol) and HOBT (0.0382g, 0.28 mmol) in 3 mL of dry DMF at 0° C, followed by the addition of EDCI (0.0541g, 0.28 mmol). Reaction was warmed up to room temperature and stirred overnight. Water was added to the reaction mixture and extracted with ethyl acetate. Organic extracts were combined, washed with saturated sodium bicarbonate, water, brine, dried over MgSO₄ and evaporated to dryness to yield a dark orange oil. Product was chromatographed on reverse phase silica using a 75:25 mixture of MeOH:H₂O, respectively to yield **35** (58.7 mg, 35% yield) as a white foam.

¹H NMR (CDCl₃, TMS) δ: 8.25 (d, 1H), 7.72 (d, 2H, *j* = 8.42 Hz), 7.50 (d, 1H), 7.24 (m, 3H), 6.95 (d, 2H, *j* = 8.42 Hz), 6.63 (d, 1H, *j* = 3.66 Hz), 5.28 (t, 1H), 5.07 (d, 2H, *j* = 9.16 Hz), 3.93 (d, 2H), 3.52 (t, 2H), 3.09 (m, 2H), 2.70 (d, 3H), 1.87-1.58 (m, 14H), 1.48-0.85 (m, 4H).


³¹P NMR (TPPO) δ: -21.2 ppm.

HPLC (60:40 CH₃CN/H₂O-0.1% solution of TFA): 8.817 min.

MS (ESI) *m/z* : 727/729 (M+Na), 705/707 (M+H).

MS high resolution (ESI) *m/z*: calculated: 705.2456; obtained: 705.2432.

-54-



Synthesis of phosphoramidate 36.

Diisopropylethyl amine (0.11 mL, 0.61 mmol) was added neat to a pre-cooled solution of phosphoramidate 26 (0.14g, 0.28 mmol), benzylamine 34 (0.08g, 0.31 mmol) and PyBOP (0.14g, 0.28 mmol) in 2 mL of dry DCM at 0 °C. Reaction was stirred for 10 min at 0 °C and 30 minutes at room temperature. Reaction was quenched with saturated ammonium chloride and DCM. Layers were separated; organic layer was washed with brine, dried over MgSO₄ and evaporated to dryness to yield light yellow oil. Crude product was chromatographed using a mixture of 9:1 EtOAc/MeOH to yield 36 (0.16g, 75% yield) as white foam.

¹H NMR (CDCl₃, TMS) δ: 8.34-8.20 (m, 2H), 8.11-8.04 (m, 1H), 7.92-7.78 (bs, 1H), 7.71 (d, 2H, *j* = 8.60 Hz), 7.63-7.43 (m, 3H), 7.28-7.24 (m, 2H), 6.95 (d, 1H, *j* = 8.6 Hz), 6.8 (s, 1H), 6.27 (d, 1H, *j* = 7.79 Hz), 5.84-5.67 (bs, 1H), 5.43-5.16 (m, 3H), 3.94 (s, 3H), 3.92 (s, 3H), 3.47 (t, 2H), 3.21-2.94 (m, 2H), 2.69 (d, 3H, *j* = 10.52 Hz), 1.96-1.49 (m, 14H), 1.41-0.93 (m, 5H).

³¹P NMR (TPPO) δ: -18.82 ppm.


HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 24.05 min.

MS (ESI) *m/z* : 802/804 (M+Na), 780/782 (M+H).

MS High Resolution (ESI) *m/z*: calculated: 780.3181; obtained: 780.3168.

25

-55-



Synthesis of prodrug 37.

Ceric ammonium nitrate (0.11g, 0.21 mmol) was dissolved in 1 mL of H₂O and added slowly to a solution of phosphoramidate 36 (0.08g, 99.83 μmol) in 3.7 mL of acetonitrile and 1.9 mL of H₂O. Reaction was stirred for 1.5 hours and then quenched by the addition of saturated ammonium chloride and extracting it with EtOAc. Organic layers were combined, washed with brine, dried over MgSO₄ and evaporated to dryness to yield light yellow oil. Crude product was chromatographed using a mixture of 9:1 EtOAc/MeOH to yield 37 (0.03g, 65% yield) as light yellow foam.

15 ¹H NMR (CDCl₃, TMS) δ : 8.34-8.21 (m, 2H), 8.11-8.04 (m, 1H), 7.92-7.78 (bs, 1H), 7.72 (d, 2H, j = 8.60 Hz), 7.28-7.24 (m, 2H), 6.95 (d, 2H, j = 8.60 Hz), 6.94 (d, 1H, j = 8.6 Hz), 6.9 (s, 1H), 6.41 (d, 1H, j = 8.6 Hz), 5.84-5.67 (bs, 1H), 5.43-5.16 (m, 3H), 3.93 (s, 3H), 3.91 (s, 5H), 3.47 (t, 2H), 3.21-2.94 (m, 2H), 2.69 (d, 3H, j = 10.52 Hz), 1.96-1.49 (m, 14H), 1.41-0.93 (m, 5H).


20 ³¹P NMR (TPPO) δ : -19.0

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 21.45 min.

MS (ESI) m/z : 772/774 (M+Na), 750/752 (M+H).

MS high resolution (ESI) m/z : calculated: 750.2711; obtained: 750.2708.

-56-




Synthesis of prodrug 38.

Prodrug 38 was synthesized in a similar manner as described for 36.

- 5 ^1H NMR (CDCl_3 , TMS) δ : 8.26-8.11 (m, 1H), 8.09-7.99 (m, 1H), 7.95-7.76 (m, 1H), 7.24-7.09 (m, 2H), 7.08-6.99 (m, 2H), 6.96-6.82 (m, 1H), 6.72-6.55 (m, 1H), 6.50-6.20 (m, 1H), 6.18-6.54 (m, 1H), 5.16-4.82 (m, 3H), 4.79-4.46 (m, 1H), 4.03-3.78 (m, 2H), 3.62-3.43 (m, 2H), 3.15-2.81 (m, 5H), 2.80-2.51 (m, 3H), 2.04-1.89 (m, 3H), 1.87-1.56 (m, 10H), 1.49-0.90 (m, 8H) ppm.
- 10 ^{31}P NMR (TPPO) δ : -18.94 ppm.
- HPLC (gradient 30-100%, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA, over 35 min): 19.03, 19.25 min.
- MS (ESI) m/z : 790/792 ($\text{M}+\text{H}$), 812/814 ($\text{M}+\text{Na}$)
- MS High Resolution (ESI) m/z : calculated: 790.2984; obtained: 790.2981.

15



Synthesis of prodrug 39.

Prodrug 39 was synthesized in a similar manner as described for 35.

- ^1H NMR (CDCl_3 , TMS) δ : 8.21-8.09 (m, 1H), 8.08-7.98 (m, 1H), 7.93-7.71 (m, 2H), 7.23-7.04 (m, 6H), 6.66-6.47 (m, 1H), 6.36-6.16 (m, 1H), 6.11-5.80 (m, 1H), 5.30-5.15

-57-

(m, 2H), 5.14-4.81 (m, 1H), 4.75-4.48 (m, 1H), 4.02-3.85 (m, 2H), 3.62-3.42 (m, 2H), 3.13-2.82 (m, 5H), 2.82-2.58 (m, 3H), 2.04-1.92 (m, 3H), 1.84-1.53 (m, 9H), 1.51-0.87 (m, 9H) ppm.


^{31}P NMR (TPPO) d: -19.16 ppm.

5 HPLC (gradient 30-100%, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA, over 35 min): 20.17, 20.35 min.

MS (ESI) m/z : 806/808 ($\text{M}+\text{H}$), 828/830 ($\text{M}+\text{Na}$)

MS High Resolution (ESI) m/z : calculated: 806.2755; obtained: 806.2758.

10



Synthesis of prodrug 40.

Prodrug 40 was synthesized in a similar manner as described for 35.

^1H NMR (CDCl_3 , TMS) d: 8.10-8.00 (m, 1H), 7.99-7.86 (m, 1H), 7.45-7.09 (m, 7H), 6.98-6.85 (m, 1H), 6.69-6.56 (m, 1H), 6.05-5.89 (m, 1H), 5.87-5.69 (m, 1H), 5.19-4.95 (m, 3H), 4.02-3.81 (m, 2H), 3.64-3.29 (m, 4H), 3.26-2.92 (m, 2H), 2.87-2.52 (m, 3H), 2.02-1.57 (m, 11H), 1.54-0.89 (m, 6H).


^{31}P NMR (TPPO) d: -18.91, -18.99 ppm.

HPLC (gradient 30-100%, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ -0.1% TFA, over 35 min): 20.65 min.

MS (ESI) m/z : 719/721 ($\text{M}+\text{H}$), 741/743 ($\text{M}+\text{Na}$)

20 MS High Resolution (ESI) m/z : calculated: 719.2613; obtained: 719.2600.

-58-



Synthesis of prodrug 41.

Prodrug 41 was synthesized in a similar manner as described for 35.


¹H NMR (CDCl₃, TMS) δ: 8.14-8.00 (m, 1H), 7.93-7.80 (m, 1H), 7.81-7.69 (m, 1H),
 5 7.42-7.30 (m, 1H), 7.27-7.06 (m, 5H), 7.04-6.93 (m, 1H), 6.93-6.81 (m, 1H), 6.23-5.99
 (m, 2H), 5.27-5.15 (m, 2H), 5.13-4.93 (m, 1H), 3.99-3.81 (m, 2H), 3.60-3.38 (m, 4H),
 3.21-2.98 (m, 2H), 2.85-2.62 (m, 3H), 1.92-1.59 (m, 11H), 1.50-0.97 (6H).

³¹P NMR (CDCl₃, TPPO) δ: -19.26

HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 21.77 min.

10 MS (ESI) *m/z* : 735/737 (M+H), 757/759 (M+Na).

MS High Resolution (ESI) *m/z*: calculated: 735.2384; obtained: 735.2358.



Synthesis of prodrug 42.

15 Prodrug 42 was synthesized in a similar manner as described for 35.

¹H NMR (CDCl₃, TMS) δ: 8.22 (d, 1H, *j* = 2.01 Hz), 7.96-7.81 9m, 1H), 7.61 (d, 2H,
j = 8.15 Hz), 7.56-7.40 (m, 1H), 7.25-7.13 (m, 1H), 6.92 (d, 2H, *j* = 8.15 Hz), 6.82-6.66
 (m, 1H), 6.66-6.53 (m, 1H), 6.27-6.00 (m, 2H), 5.35-5.10 (m, 1H), 5.03 (d, 2H, *j* = 9.25
 Hz), 3.90 (d, 2H, *j* = 5.86 Hz), 3.48 (t, 2H), 3.18-2.92 (m, 2H), 2.64 (d, 3H, *j* = 10.53 Hz),
 20 1.94-1.47 (m, 10H), 1.38-0.98 (m, 7H) ppm.

-59-

³¹P NMR (CDCl₃, TPPO) δ: -14.96 ppm.HPLC (gradient 30-100%, CH₃CN/H₂O-0.1% TFA, over 35 min): 17.28 min.MS (ESI) *m/z* : 704/706 (M+H), 726/728 (M+Na).MS High Resolution (ESI) *m/z*: calculated: 704.2616; obtained: 704.2633.

5

Growth Inhibition in L1210 Mouse Leukemia Cells

Stock solutions of drugs were prepared in absolute ethanol, and serial dilutions of drug were prepared in ethanol. L1210 cells were suspended in Fisher's medium supplemented with 10% horse serum, 1% glutamine, and 1% antibiotic-antimycotic solution to give 10 mL volumes of cell suspension at a final density of 3-6 X 10⁴ cells/mL. Appropriate volumes of the drug solution were transferred to the cell suspensions, and incubation was continued for 2, 8, 24, or 48 hours. The cells were spun down, resuspended in fresh drug-free medium and returned to the incubator. Final cell counts were determined 48 hours after the start of drug treatment. The data were analyzed by sigmoidal curve fit of cell count vs. Log (drug concentration) and the results expressed as the IC₅₀ (the drug concentration that inhibits cell growth to 50% of control value).

20

Compound	IC ₅₀ nM			
	2h	8h	24h	48h
3	116	47	7.2	2.3
5	44	16	3.1	.63
6	34	48	4.8	1.6
5-FU	2500	1010	360	200
FUdR	45	23	4.1	.64

25

Luciferase inhibition on transfected J77 cells.

J77 cells (2 X 10⁷ cells in 500 mL of FBS-free RPMI media) were transfected with 15 mg of NF-AT-Luciferase plasmid. Cells were transferred to a flask with 10 mL of complete RPMI media and incubated for 24 hours. After this time cells were spun down, resuspended in 12 mL of complete RPMI media and divided into a 6-

30

-60-

well plate (2 mL/well). Drug stock solutions in DMSO were made and each well was treated with a different concentration and incubated for 2 hours. Each well was then treated with 10 mL of a mixture of 1.25 mg of PMA and 9 mg of Ionomycin in 115 mL of FBS-free RPMI, followed by the addition of 2 mg of Anti-CD3. Cells were incubated
5 for 6 hours, spun down, washed with PBS, lysed for 15 min and centrifuged. Supernatant was collected and stored at -78°C overnight. Luciferase activity was measured using a luminometer with a mixture of 50 mL of luciferase substrate and 10 mL of supernatant.

10	Compound	IC₅₀ (uM)
	35	34.1
	37	4.2
	38	5.6
	40	3.4
15	42	15.0

Tritium Assay in L1210, LM and LM(TK-) Cells

Stock solutions of drugs were prepared in absolute ethanol. L1210 cells
20 were suspended in fisher's medium, LM and LM (TK-) cells were suspended in minimum essential medium to give a final density of 11.24×10^6 cells/mL. Aliquots (445 mL) of the cell suspension were placed in ependorf tubes, followed by addition of 5 mL of drug solution and incubated at 37°C for 2 hours. Tritium release reaction was started by the addition of 50 mL of serum free media containing 1 mCi $[5\text{-}^3\text{H}]\text{dCyt}$ in a
25 0.5 mM dCyt solution. Allowed reaction to incubate for 60 min in a shaking water bath at 37°C . Reaction was terminated by transferring 100 mL of the reaction mixture to ependorf tubes containing 100 mL of a 20% mixture of activated charcoal suspension in 4% aqueous perchloric acid. Tubes were vigorously vortexed, then centrifuged in a microfuge. Radioactivity of 100 mL of the supernatant was measured using a
30 scintillation spectrometer. The data were analyzed by sigmoidal curve fit of CMP vs. Log (drug concentration) and the results expressed as the IC₅₀ (the drug concentration that inhibits cell growth to 50% of control value).

-61-

Compound	IC ₅₀ nM		
	L1210	LM	LM(TK-)
3	46	711	1280
FudR	8	59	3670

5

Pharmaceutical Formulations

The following parenteral formulations are prepared for use in cancer therapy.

10

<u>Compound</u>	<u>Mass/Dose (mg)</u>	<u>Carrier(s)</u>
3	25	isotonic saline
5	50	isotonic saline + DMSO
6	100	5% glucose
15 35	550	5% glucose + cyclodextrin
37	180	
38	300	lipid emulsion
40	40	isotonic saline
20 42	450	isotonic saline + cyclodextrin

25

-62-

CLAIMS:

1. A compound of the formula

$$R_rCH_2OP(halo)NR(CH_2)_nX$$
 - 5 wherein
 - R is C₁-C₄ alkyl or -(CH₂)_nX;
 - n is 4 or 5;
 - X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;
 - 10 halo is chloro, bromo or iodo; and
 the group R_rCH₂- is a biologically labile ester forming group.
2. The compound of claim 1 wherein n is 4.
3. The compound of claim 1 wherein n is 5.
4. The compound of claim 1 wherein R is methyl.
- 15 5. The compound of claim 1 wherein halo is chloro.
6. The compound of claim 1 wherein X is chloro or bromo.
7. A method of preparing a phosphoramidate prodrug for enhanced intracellular delivery of a drug as phosphate ester or amide said method comprising the steps of reacting a hydroxy functional or amino functional drug compound (Drug-ZH)
 - 20 with a compound of the formula

$$R_rCH_2OP(halo)NR(CH_2)_nX$$
 under conditions conducive to the formation of an intermediate compound of the formula

$$R_rCH_2OP(Z-Drug)NR(CH_2)_nX$$
 and oxidizing that intermediate to form the phosphoramidate prodrug of the formula

$$R_rCH_2OP(O)(Z-Drug)NR(CH_2)_nX$$
 - 25 in which formulas
 - R is C₁-C₄ alkyl or -(CH₂)_nX;
 - n is 4 or 5;
 - Z is O or N;
 - 30 X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;
 - halo is chloro, bromo or iodo; and

-63-

the group R_1CH_2- is a biologically labile ester forming group.

8. The method of claim 7 wherein Drug-ZH is an amino acid, or a biologically active peptide or peptidomimetic.

9. The method of claim 8 wherein Drug-ZH is a peptidomimetic of
5 the formula



10

wherein Z is O or N;

q and k are independently 1 or 0; and

B is H, amino, protected amino or C₁-C₄ alkanoylamino.

10. The method of claim 7 wherein Drug-ZH is a biologically active
15 nucleotide analog.

11. A phosphoramidate compound formed from a hydroxy functional or amino functional drug compound of the general formula Drug-ZH said prodrug being a compound of the formula



20 wherein

R is C₁-C₄ alkyl or -(CH₂)_nX;

n is 4 or 5;

Z is O or N:

25 X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;

halo is chloro, bromo or iodo; and

the group R_1CH_2- is a biologically labile ester forming group.

12. The prodrug of claim 11 wherein the drug is an amino acid, or a biologically active peptide or peptidomimetic.

30 13. The method of claim 12 wherein Drug-ZH is a peptidomimetic of
the formula

-64-



5

wherein Z is O or N;

q and k are independently 1 or 0; and


B is H, amino, protected amino or C₁-C₄ alkanoylamino.

14. The prodrug of claim 11 wherein the drug is a biologically active
10 nucleotide analog.

15. A method of preparing a compound of the formula



comprising the steps of reacting a compound of the formula



- 15 with 1) an alcohol of the formula R₁CH₂OH and 2) an amine of the formula
HNR(CH₂)_nX, each in the presence of an acid scavenger,

wherein in the above formulas

m is 0 or 1;

R is C₁-C₄ alkyl or -(CH₂)_nX;

- 20 n is 4 or 5;

X is an electrophilic group capable of being nucleophilically displaced
from its bonded carbon atom;

halo is chloro, bromo or iodo; and

the group R₁CH₂- is a biologically labile ester forming group.

- 25 16. A method of preparing a phosphoramidate prodrug of the formula



for enhanced intracellular delivery of a compound of the general formula Drug-ZPO₃
said method comprising the steps of reacting a hydroxy functional amino functional drug
compound of the formula Drug-ZH with a compound of the formula

- 30 $R_1CH_2OP(O)(halo)NR(CH_2)_nX$

under conditions conducive to the formation of the prodrug

wherein in the above formulas

-65-

R is C₁-C alkyl or -(CH₂)_nX;

n is 4 or 5;

Z is O or N;

X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;

halo is chloro, bromo or iodo; and

the group $R\text{CH}_2-$ is a biologically labile ester forming group.

17. A pharmaceutical composition comprising

a phosphoramidate compound formed from a hydroxy functional or amino

10 functional drug compound of the general formula Drug-ZH said prodrug being a
compound of the formula



wherein

R is C₁-C₄ alkyl or -(CH₂)_nX;

n is 4 or 5:

Z is O or N:

X is an electrophilic group capable of being nucleophilically displaced from its bonded carbon atom;

halo is chloro, bromo or iodo;

the group R_1CH_2- is a biologically labile ester forming group; and a pharmaceutically acceptable carrier therefor.

18. The pharmaceutical compound of claim 17 wherein Drug-ZH is an amino acid or a biologically active peptide or peptidomimetic.

19. The pharmaceutical composition of claim 18 wherein Drug-ZH is
25 a peptidomimetic of the formula



wherein Z is O or N:

q and k are independently 1 or 0; and

-66-

B is H, amino, protected amino or C₁-C₄ alkanoylamino.

20. The pharmaceutical composition of claim 17 wherein Drug-ZH is a nucleotide analog.

INTERNATIONAL SEARCH REPORT

ational Application No
PCT/US 01/10163

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07F9/26 A61K31/66 A61K31/70 A61P35/00 C07F9/655
C07F9/6553 C07F9/24 C07H19/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07F C07H A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, BEILSTEIN Data, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEN R.Y. ET AL.: "The synthesis of 0-monosaccharidyl-methoxycarbonyl-phosphonamides by Arbuzov reaction" CHINESE CHEMICAL LETTERS., vol. 6, no. 1, 1995, pages 23-26, XP001010603 XX, XX ISSN: 1001-8417 page 24	1
A	WO 93 06120 A (UNIVERSITY OF ROCHESTER) 1 April 1993 (1993-04-01) the whole document --- -/-	11-14, 17-20

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

31 July 2001

Date of mailing of the international search report

09/08/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx: 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/10163

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SIDDIQUI A Q ET AL: "DESIGN AND SYNTHESIS OF LIPOPHILIC PHOSPHORAMIDATE D4T-MP PRODRUGS EXPRESSING HIGH POTENCY AGAINST HIV IN CELL CULTURE: STRUCTURAL DETERMINANTS FOR IN VITRO ACTIVITY AND QSAR" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 42, no. 20, 1999, pages 4122-4128, XP000999173 ISSN: 0022-2623 the whole document	11-14, 17-20
A	MCGUIGAN C ET AL: "PHOSPHORAMIDATES AS POTENT PRODRUGS OF ANTI-HIV NUCLEOTIDES: STUDIES IN THE AMINO REGION" ANTIVIRAL CHEMISTRY & CHEMOTHERAPY, GB, BLACKWELL SCIENTIFIC PUBL., LONDON, vol. 7, no. 1, 1996, pages 31-36, XP000578787 ISSN: 0956-3202 the whole document	11-14, 17-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/10163

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9306120 A	01-04-1993	US 5233031 A	03-08-1993
		CA 2119351 A	01-04-1993
		EP 0605582 A	13-07-1994
		JP 6511003 T	08-12-1994